



INSTITUT
POLYTECHNIQUE
DE PARIS

Journal de l'École polytechnique

Mathématiques

Rafik IMEKRAZ

A necessary and sufficient condition for probabilistic continuity on a boundaryless compact Riemannian manifold

Tome 9 (2022), p. 747-805.

http://jep.centre-mersenne.org/item/JEP_2022__9__747_0

© Les auteurs, 2022.

Certains droits réservés.



Cet article est mis à disposition selon les termes de la licence
LICENCE INTERNATIONALE D'ATTRIBUTION CREATIVE COMMONS BY 4.0.
<https://creativecommons.org/licenses/by/4.0/>

L'accès aux articles de la revue « Journal de l'École polytechnique — Mathématiques » (<http://jep.centre-mersenne.org/>), implique l'accord avec les conditions générales d'utilisation (<http://jep.centre-mersenne.org/legal/>).

Publié avec le soutien
du Centre National de la Recherche Scientifique



Publication membre du
Centre Mersenne pour l'édition scientifique ouverte
www.centre-mersenne.org

A NECESSARY AND SUFFICIENT CONDITION FOR PROBABILISTIC CONTINUITY ON A BOUNDARYLESS COMPACT RIEMANNIAN MANIFOLD

BY RAFIK IMEKRAZ

ABSTRACT. — We give a necessary and sufficient condition for the uniform convergence of random series of eigenfunctions on a boundaryless compact Riemannian manifold. As a consequence, we generalize an estimate of Burq and Lebeau about the supremum of a random eigenfunction. Finally, we prove that our results are universal with respect to the random variables (this is a Riemannian analogue of a result of Marcus and Pisier), with respect to compact submanifolds and with respect to the Riemannian structure of the manifold. Our proofs rely on several tools like the Dudley-Fernique theorem, the Slepian comparison theorem and the semi-classical functional calculus for elliptic operators on compact manifolds.

RÉSUMÉ (Condition nécessaire et suffisante pour la continuité probabiliste sur une variété riemannienne compacte sans bord)

Nous donnons une condition nécessaire et suffisante pour la convergence uniforme de séries aléatoires de fonctions propres sur une variété riemannienne compacte sans bord. Comme conséquence, nous généralisons une estimation de Burq et Lebeau concernant les bornes d'une fonction propre aléatoire. Finalement, nous prouvons que nos résultats sont universels par rapport aux variables aléatoires utilisées (il s'agit d'un analogue riemannien d'un résultat de Marcus et Pisier), par rapport aux sous-variétés compactes et à la structure riemannienne de la variété compacte. Nos preuves reposent sur plusieurs outils dont le théorème de Dudley-Fernique, le théorème de comparaison de Slepian et un calcul fonctionnel semi-classique pour les opérateurs elliptiques d'une variété compacte.

CONTENTS

1. Introduction and statement of the main result.....	748
2. Statements of the main results in the Gaussian case.....	757
3. Application to universality results for random series.....	759
4. Metric considerations.....	761
5. Partial pseudo-distances on spheres.....	764
6. Partial pseudo-distances on manifolds.....	768
7. Preliminaries for proofs of Theorems 2 and 3.....	775
8. Proof of Theorem 2 for spheres S^d with $d > 2$	777
9. Proof of Theorem 3 for S^d with $d > 2$	783

MATHEMATICAL SUBJECT CLASSIFICATION (2020). — 60G50, 60G15, 58J40, 46B09.

KEYWORDS. — Paley-Zygmund theorems, Laplace-Beltrami operator, compact manifolds, Gaussian processes.

Partially supported by the ANR projects ESSED ANR-18-CE40-0028 and UNIRANDOM ANR-17-CE40-0008.

10. Proof of Theorem 2 and 3 for compact manifolds.....	784
11. Proof of (18) in Theorem 1, quantitative version.....	785
12. Proof of Theorem 1, qualitative version.....	790
13. Proof of Theorem 5, semi-classical analysis and independence of the Riemannian metric.....	792
Appendix A. First moments of chi distributions.....	799
Appendix B. Proof of Theorem 11 via a result of Canzani-Hanin.....	800
References.....	803

1. INTRODUCTION AND STATEMENT OF THE MAIN RESULT

Let \mathcal{M} be a boundaryless compact Riemannian manifold of dimension $d > 2$, the goal of Theorem 1 is to give a necessary and sufficient condition to ensure the convergence, in the Banach space $\mathcal{C}^0(\mathcal{M})$, of suitable random linear combinations of eigenfunctions on \mathcal{M} . The problem studied is posed in the paper of Tzvetkov [Tzv10, p. XV-6] and, as it will be explained in this introduction, is equivalent to control expectations of supremum of linear combinations of eigenfunctions. Thus we need to generalize an optimal two-sided inequality obtained by Burq and Lebeau in [BL13, BL14], see (15) and the new inequalities (16) and (18). The results below are Riemannian analogues of those of the classical theory, for \mathcal{M} being a compact group, whose final treatment is done in the book of Marcus and Pisier [MP81]. And we moreover prove, via the semi-classical analysis, that all our results are independent of the Riemannian structure of \mathcal{M} (see Theorem 5).

Our result really looks like a sufficient condition obtained by Salem and Zygmund in [SZ54, p. 291], the main new point is that our Salem-Zygmund condition is necessary and sufficient in the manifold framework (see (19) in Theorem 1) whereas it is merely sufficient (and not necessary) in the classical theory on the torus (see [SZ54, p. 292]). As underlined in the paper [BL13], a clue suggesting that better results should exist in the Riemannian setting is that the dimensions of the eigenspaces of the Laplace-Beltrami operator on S^d , for $d > 2$, tend to $+\infty$ in contrast to the classical setting $L^2(\mathbb{T}) = \bigoplus_{n \in \mathbb{Z}} \mathbb{C}e^{in \cdot}$. The present contribution gives another enlightenment (see below Theorem 1 for definitions): it turns out that the Dudley pseudo-distance of the Gaussian random wave of each eigenspace of S^d is equivalent to a very simple distance (see Theorems 8 and 10).

Before going into details on compact manifolds (and giving a precise definition of our random series), it is worthwhile to recall the main results on the torus $\mathbb{T} = \mathbb{R}/2\pi\mathbb{Z}$ (we refer to [MP81] for the general case with compact groups). Although the first results had been proved for Fourier series with real trigonometric functions, it is known that there is no difference to deal with complex trigonometric functions (see [MP81, p. 122]). For any real sequence $(c_n)_{n \in \mathbb{N}} \in \ell^2(\mathbb{N})$, let us consider the function $f \in L^2(\mathbb{T})$ defined by

$$f(x) = \sum_{n \in \mathbb{N}} c_n e^{inx}.$$

Now fix a sequence $(\varepsilon_n)_{n \in \mathbb{N}}$ of i.i.d. Rademacher random variables, in other words $\mathbf{P}(\varepsilon_n = 1) = \mathbf{P}(\varepsilon_n = -1) = 1/2$. With probability 1 with respect to ω running over a reference probability space Ω , Paley and Zygmund proved in [PZ30, p. 347] that the following random series

$$(1) \quad f^\omega(x) := \sum_{n \in \mathbb{N}} \varepsilon_n(\omega) c_n e^{inx}$$

almost surely converges in $\mathcal{C}^0(\mathbb{T})$ provided that the following condition is fulfilled

$$(2) \quad \exists \gamma > 1, \quad \sum_{n=2}^{+\infty} c_n^2 \ln^\gamma(n) < +\infty.$$

The strong motivation of the resurgence of conditions like (2) is that they can be interpreted as probabilistic Sobolev embeddings (more details are given in the introductions of [Tzv10, BL13] and [Ime19, p. 2734]). In [SZ54, p. 291], Salem and Zygmund relaxed the assumption (2) by the following one

$$(3) \quad \sum_{n=2}^{+\infty} \frac{1}{n \sqrt{\ln(n)}} \left(\sum_{k=n}^{+\infty} c_k^2 \right)^{1/2} < +\infty,$$

and also remarked that such a condition is not necessary. An important step is made by Marcus in the papers [Mar73, Mar75] in which we learn that (3) becomes a necessary and sufficient condition if the sequence $(|c_n|)_{n \in \mathbb{N}}$ in (1) is non-increasing and if the Rademacher random variables ε_n are replaced with a sequence $(g_n^{\mathbb{C}})_{n \in \mathbb{N}}$ of independent complex standard Gaussian random variables⁽¹⁾ $\mathcal{N}_{\mathbb{C}}(0, 1)$ (see [Mar75] for general complex symmetric random variables satisfying suitable normalization conditions).

For the problem of finding a necessary and sufficient condition, the solution had finally come from a drastically different point of view. More precisely, for any sequence $(c_n) \in \ell^2(\mathbb{Z})$, the general Gaussian Fourier series

$$(4) \quad f^{G,\omega}(x) = \sum_{n \in \mathbb{Z}} g_n^{\mathbb{C}}(\omega) c_n e^{inx}$$

is now seen as a stationary Gaussian random process on the torus \mathbb{T} . Here the word “stationary” means that the random process $(f^G(x))_{x \in \mathbb{T}}$ is invariant under the group action of \mathbb{T} : in particular, for any angle $\alpha \in \mathbb{R}$, $f^{G,\omega}(x + \alpha)$ has the same distribution as that of $f^{G,\omega}(x)$ due to the complex symmetry of the complex Gaussian variables $g_n^{\mathbb{C}}$. It is worthwhile to underline that such a stationary assumption has no sense for a general compact manifold \mathcal{M} . Then the important result, now called the Dudley-Fernique theorem, allows for proving that the almost sure continuity of the Gaussian functions $x \mapsto f^{G,\omega}(x)$ is equivalent to the so-called entropy condition:

$$(5) \quad \int_0^{+\infty} \sqrt{\ln N_\delta(\mathbb{T}, \varepsilon)} d\varepsilon < +\infty,$$

⁽¹⁾In particular we have $\mathbf{E}[g_n^{\mathbb{C}}] = 0$ and $\mathbf{E}[|g_n^{\mathbb{C}}|^2] = 1$.

where

- the function $\delta : \mathbb{T} \times \mathbb{T} \rightarrow \mathbb{R}^+$ is the Dudley pseudo-distance of \mathbb{T} defined by

$$\delta(x, y)^2 = \mathbf{E} [|f^{G, \omega}(x) - f^{G, \omega}(y)|^2] = 4 \sum_{n \in \mathbb{Z}} |c_n|^2 \sin^2 \left(\frac{n(x-y)}{2} \right),$$

- the number $N_\delta(\mathbb{T}, \varepsilon)$ is the covering number of the torus \mathbb{T} with respect to δ , namely the minimal number of open δ -balls of radius $\varepsilon > 0$ whose union covers \mathbb{T} .

Moreover, Marcus and Pisier proved that one may replace the sequence of Gaussian random variables in (4) with any sequence of independent real centered random variables $(X_n)_{n \in \mathbb{Z}}$ satisfying

$$0 < \inf_{n \in \mathbb{Z}} \mathbf{E}[|X_n|] \quad \text{and} \quad \sup_{n \in \mathbb{Z}} \mathbf{E}[|X_n|^2] < +\infty.$$

Such a result is usually called a universality phenomenon and allows for coming back to the Rademacher random variables $X_n = \varepsilon_n$ (see [MP81, p. 7–9] and [Pis78, p. 28, Cor. 7.3]).

One may think that the entropy condition (5) is a bit abstract but it really captures the essence of the almost sure continuity. Firstly, the entropy condition allows for showing the sufficiency of the more concrete Paley-Zygmund (2) and Salem-Zygmund (3) assumptions (see [MP81, Part VII.1]). Secondly, the entropy condition (which is of qualitative nature) is equivalent to the following quantitative version (see [MP81, p. 11] or [Pis78, p. 3–4]):

$$(6) \quad \mathbf{E}_\omega \left[\left\| \sum_{k=-n}^n \varepsilon_k(\omega) c_k e^{ikx} \right\|_{\mathcal{C}_x^0(\mathbb{T})} \right] \simeq \mathbf{E}_\omega \left[\left\| \sum_{k=-n}^n g_k^{\mathbb{C}}(\omega) c_k e^{ikx} \right\|_{\mathcal{C}_x^0(\mathbb{T})} \right] \\ \simeq |c_0| + \int_0^{+\infty} \sqrt{\ln N_\delta(\mathbb{T}, \varepsilon)} d\varepsilon.$$

For instance, here are the so-called Salem-Zygmund inequalities (see [LQ18b, p. 259]):

$$(7) \quad \forall n \gg 1, \quad \mathbf{E}_\omega \left[\frac{1}{\sqrt{n}} \left\| \sum_{k=n+1}^{2n} \varepsilon_k(\omega) e^{ikx} \right\|_{\mathcal{C}_x^0(\mathbb{T})} \right] \simeq \sqrt{\ln(n)}.$$

For the sequel, it is worthwhile to note that the Dudley pseudo-distance δ_n of

$$\frac{1}{\sqrt{n}} \sum_{k=n+1}^{2n} g_k^{\mathbb{C}}(\omega) e^{ikx}$$

satisfies the following estimates⁽²⁾

$$(8) \quad \forall n \gg 1, \quad \delta_n \simeq \min(1, n\delta_g), \quad \text{where } \delta_g \text{ is the Riemannian distance of } \mathbb{T},$$

which in turn allows to show the implication (6) \Rightarrow (7).

⁽²⁾By setting $\theta = (x-y)/2$, one checks that (8) comes from the exact formulas $\delta_n(x, y)^2 = (4/n) \sum_{k=n+1}^{2n} \sin^2(k\theta) = 2 - (2 \sin(n\theta)/n \sin(\theta)) \cos((3n+1)\theta)$ by separating the cases $0 \leq \theta \leq 2/n$ and $2/n \leq \theta \leq \pi/2$.

It is time to recall the known literature about the generalization of the previous results to a boundaryless compact Riemannian manifold \mathcal{M} of dimension $d > 2$. We denote by Δ the non-positive Laplace-Beltrami operator of \mathcal{M} . The Hilbert space $L^2(\mathcal{M})$ is considered with respect to the Riemannian measure $\text{vol}_{\mathcal{M}}$ of \mathcal{M} . We recall that $L^2(\mathcal{M})$ admits a Hilbert basis $(\phi_k)_{k \in \mathbb{N}}$ of eigenfunctions of Δ :

$$(9) \quad \Delta \phi_k = -\lambda_k^2 \phi_k, \quad 0 = \lambda_0 < \lambda_1 \leq \lambda_2 \leq \dots \rightarrow +\infty.$$

For simplicity, we shall assume that each ϕ_k is real-valued. By fixing a sequence of coefficients $(c_k)_{k \in \mathbb{N}}$, a natural option is to replace the random Fourier series (1) and (4) with the following random series:

$$(10) \quad \sum_{k \in \mathbb{N}} \varepsilon_k(\omega) c_k \phi_k(x) \quad \text{or} \quad \sum_{k \in \mathbb{N}} g_k(\omega) c_k \phi_k(x), \quad (\omega, x) \in \Omega \times \mathcal{M},$$

where $(g_k)_{k \in \mathbb{N}}$ is a sequence of i.i.d. real Gaussian random variables $\mathcal{N}_{\mathbb{R}}(0, 1)$. For the random series (10), Tzvetkov generalized in [Tzv10] the Paley-Zygmund condition (2), that is a sufficient condition ensuring the almost sure convergence in $\mathcal{C}^0(\mathcal{M})$. In contrast with the sequence of eigenvalues $(\lambda_k^2)_{k > 0}$, the sequence of eigenfunctions $(\phi_k)_{k > 0}$ may not be unique (actually if Δ has multidimensional eigenspaces). For instance, on $\mathcal{M} = S^d$, there are sub-sequences of eigenfunctions having very different behaviors like concentration around a point or around a geodesic (those are called zonal eigenfunctions or Gaussian beams). Without any further information on the sequence $(\phi_k)_{k \in \mathbb{N}}$, it seems hopeless to expect to have a simple, necessary and sufficient condition for the almost sure convergence in $\mathcal{C}^0(\mathcal{M})$ of (10). By comparison with the classical theory on a compact group G , the adequate random series are defined by taking account of the irreducible representations of G (see [FTR67, MP81]). In the Riemannian framework, it is thus natural to make an additional restriction that counterbalances the possible concentration of sub-sequences of eigenfunctions. The option chosen by Burq and Lebeau in [BL13] will be detailed below and relies in gathering eigenfunctions whose eigenvalues are in a same suitable small interval (for instance for the same eigenvalue for $\mathcal{M} = S^d$, see (11)). We also refer to [Ime18, p.272] and also [IRT16, Ime19] for more details about that multidimensional point of view.

Before writing the slight modification of (10), let us underline that the situation is drastically different for the L^p case with $p < +\infty$. Indeed, there is a very nice formula of Maurey that gives a necessary and sufficient condition for the almost sure convergence of unidimensional series like (1) and (10) in L^p for $p < +\infty$ (see [Mau74, p.22, Cor.1], [LT79, Th.1.d.6] or [IRT16, Prop.2.1] for a recent use). For a multidimensional analogue of the Maurey formula, we refer to [Ime18, Th.2.1 & 2.21] and [Ime19, Th.12.1]. Without going into details on the difficulties of the L^p case for $p < +\infty$, one could say that the analogue problems of finding necessary and sufficient conditions on L^p are quite well understood for finite p (see the papers [AT08, Tzv10, Gri10, IRT16, Ime18] and [Ime19, Th.2.3, 4.5 & 4.6]).

It is time to properly define the random series that naturally replace (10) in the case of the sphere S^d (the case of manifold is considered just after). We recall that the

sequence of eigenvalues of the Laplace-Beltrami Δ on the sphere S^d is given, without counting multiplicities, by $-n(n+d-1)$ with $n \in \mathbb{N}$. Moreover, the dimension of the eigenspace

$$(11) \quad E_n = \ker(\Delta + n(n+d-1)) \subset L^2(S^d)$$

satisfies the numerical equivalence $\dim(E_n) \simeq n^{d-1}$ for $n > 1$. For any function $f \in L^2(S^d)$, we write

$$f = \sum_{n \in \mathbb{N}} f_n \quad \text{with} \quad f_n = \sum_{\lambda_k^2 = n(n+d-1)} \langle f, \phi_k \rangle \phi_k \in E_n.$$

For any $n \in \mathbb{N}^*$, we now consider a uniform random vector $U_n : \Omega \rightarrow \mathbb{R}^{\dim(E_n)}$, namely whose probability distribution is the probability spherical measure on the unit sphere $\mathbb{S}^{\dim(E_n)-1}$. We write the coordinates $U_n = (U_{n,k})_k$ with k running over the finite set of integers such that $\lambda_k^2 = n(n+d-1)$. A natural generalization of (1) and the first random series in (10) is given by

$$(12) \quad f^\omega(x) = \sum_{n > 1} f_n^\omega(x), \quad \text{with} \quad f_n^\omega = \|f_n\|_{L^2(S^d)} \sum_{\lambda_k^2 = n(n+d-1)} U_{n,k}(\omega) \phi_k \in E_n,$$

where the uniform random vectors U_n are assumed to be mutually independent. The random function f_n^ω is thus a random eigenfunction of norm $\|f_n\|_{L^2(S^d)}$. That formalism is connected to that used by Burq-Lebeau in [BL13, App. C] (see [Ime18, p. 274] for more details).

On a general boundaryless compact Riemannian manifold \mathcal{M} , the idea is to replace the eigenspace $E_n \subset L^2(S^d)$ with the subspace $E_{(Kn-K, Kn]}$ of $L^2(\mathcal{M})$ defined as follows

$$(13) \quad E_{(Kn-K, Kn]} := \text{Span}\{\phi_k, \lambda_k \in (Kn-K, Kn]\},$$


where the spectral parameter $K > 0$ is large enough. In particular, it is known that the analogue of the asymptotic $\dim(E_n) \simeq n^{d-1}$ is given by $\dim(E_{(Kn-K, Kn]}) \simeq n^{d-1}$ (see [BL13, p. 923] or [Ime19, Lem. 8.1]). In other words, for any $f = \sum_{n > 1} f_n \in L^2(\mathcal{M})$ with $f_n \in E_{(Kn-K, Kn]}$, we set the following random series

$$(14) \quad f^\omega = \sum_{n > 1} f_n^\omega, \quad \text{with} \quad f_n^\omega := \|f_n\|_{L^2(\mathcal{M})} \sum_{\lambda_k \in (Kn-K, Kn]} U_{n,k}(\omega) \phi_k,$$

where $U_n : \Omega \rightarrow \mathbb{R}^{\dim(E_{(Kn-K, Kn]})}$ is a uniform random vector as above. At this stage of this introduction, we must recall an optimal result proved by Burq and Lebeau. By using the previous notations, [BL13, Th. 5, p. 930] states the following optimal bound on a uniform random eigenfunction f_n^ω on $L^2(S^d)$ with eigenvalue $-n(n+d-1)$:

$$(15) \quad \forall n \gg 1, \quad \mathbf{E}_\omega \left[\|f_n^\omega\|_{C^0(S^d)} \right] \simeq \sqrt{\ln(n)} \|f_n\|_{L^2(S^d)}.$$

We also refer to [CH15a, Th. 3] for a geometric control of the upper bound in (15). The idea we keep in mind is that the inequalities (15) of Burq-Lebeau should really be seen as a Riemannian analogue of the Salem-Zygmund inequalities (7).



Before stating our main result, we recall the striking equivalence of the following two statements:

- (i) for almost every $\omega \in \Omega$, the random series $\sum f_n^\omega$ converges in $\mathcal{C}^0(S^d)$,
- (ii) the random series $\sum f_n^\omega$ converges in $L^1(\Omega, \mathcal{C}^0(S^d))$.

For unidimensional Rademacher random series $\sum \varepsilon_n(\omega) f_n$, an analogue equivalence of the previous assertions was proved by Kahane and is indeed completely independent of the Banach space $\mathcal{C}^0(S^d)$ (see the reference book [Kah85] about random series, [LQ18a, p. 142, Rem. 1] or [MP81, p. 43]). For the multidimensional case we are interested in, such an equivalence is a consequence of a result by Marcus and Pisier [MP81, p. 92]. We now understand that a general solution of our problem is equivalent to find optimal bounds of the expectations $\mathbf{E}_\omega [\|\sum_{n=1}^N f_n^\omega\|_{\mathcal{C}^0(S^d)}]$ generalizing the Burq-Lebeau asymptotics (15). Our work will show the following new result on S^d (with $d > 2$):

$$(16) \quad \mathbf{E}_\omega \left[\left\| \sum_{n=1}^N f_n^\omega \right\|_{\mathcal{C}^0(S^d)} \right] \simeq \sum_{p=1}^N \frac{1}{p\sqrt{\ln(p+1)}} \left(\sum_{n=p}^N \|f_n\|_{L^2(S^d)}^2 \right)^{1/2}.$$

The right-hand side clearly looks like (3) but no monotonicity assumption is needed (in contrast to [Mar75] for the torus T). More generally, Theorem 1 settles the general case of a compact manifold with a Salem-Zygmund type condition (19) which is necessary and sufficient in contrast to the classical results (for which analogue estimates of (19) are sufficient but not necessary).

THEOREM 1. — *There is a constant $K_0 > 0$ depending only on the Riemannian manifold \mathcal{M} such that, for any $K > K_0$, if one considers*

- a sequence $(f_n)_{n>1}$ satisfying $f_n \in E_{(K_{n-K}, K_n]}$ (see (13)) for each $n \in \mathbb{N}^*$,
- a non-zero dimensional compact submanifold $\mathcal{M}_s \subset \mathcal{M}$ with smooth (eventually empty) boundary,
- a sequence of independent real random variables⁽³⁾ $(X_n)_{n>1}$ satisfying

$$(17) \quad 0 < \inf_{n>1} \mathbf{E}[|X_n|] \quad \text{and} \quad \sup_{n>1} \mathbf{E}[|X_n|^2] < +\infty,$$

⁽³⁾Note that we can choose $X_n = 1$ for each $n > 1$ in order to get (16).

then

(1) we have the numerical equivalence for any $N \in \mathbb{N}^*$:

$$(18) \quad \mathbf{E}_\omega \left[\sup_{x \in \mathcal{M}_s} \left| \sum_{n=1}^N X_n(\omega) f_n^\omega(x) \right| \right] \simeq \sum_{p=1}^N \frac{1}{p\sqrt{\ln(p+1)}} \left(\sum_{n=p}^N \|f_n\|_{L^2(\mathcal{M})}^2 \right)^{1/2},$$

where the constants of that equivalence are independent of N and depend on \mathcal{M} , \mathcal{M}_s , K , $\inf_{n>1} \mathbf{E}[|X_n|]$ and $\sup_{n>1} \mathbf{E}[|X_n|^2]$ and where we assume that all the random variables $\omega \mapsto X_n(\omega)$ and $\omega \mapsto f_n^\omega$ (defined in (14)) are mutually independent.

(2) Moreover, the following two statements are equivalent:

- the random series $\sum X_n(\omega) f_n^\omega$ almost surely converges in $\mathcal{C}^0(\mathcal{M}_s)$,
- the following Salem-Zygmund condition is fulfilled:

$$(19) \quad \sum_{p=1}^{+\infty} \frac{1}{p\sqrt{\ln(p+1)}} \left(\sum_{n=p}^{+\infty} \|f_n\|_{L^2(\mathcal{M})}^2 \right)^{1/2} < +\infty.$$

(3) Finally, the same conclusion also holds true for the particular case $\mathcal{M} = S^d$ if each f_n belongs to $\ker(\Delta + n(n+d-1))$ provided that we consider (12) instead of (14) (and thus K is irrelevant in the equivalence (18)).

Here are two comments about the previous result:

- as a consequence of Point 2) of Theorem 1, one sees that the almost sure convergence in $\mathcal{C}^0(\mathcal{M})$ is equivalent to the almost sure convergence in $\mathcal{C}^0(\mathcal{M}_s)$ for any submanifold \mathcal{M}_s (more comments in that direction are given in Section 3),
- Theorem 1 supersedes [Ime19, Th. 2.1] in which one may find a sufficient condition in the spirit of the original Paley-Zygmund theorem.

Similarly to the theory developed by Marcus and Pisier, the good idea is first to consider our problem with a suitable Gaussian analogue of $\sum f_n^\omega$. A known intuition suggests that the uniform random vector $U_n : \Omega \rightarrow S^{\dim(E_n)-1}$, in (12), is closely related to the Gaussian vector of $\mathbb{R}^{\dim(E_n)}$ with distribution $\mathcal{N}(0, 1/(\dim(E_n)) \text{Id})$ (see [Pis89, p. 58]). We then introduce the following Gaussian analogue of (12) for any $f \in L^2(S^d)$

$$(20) \quad f^{G,\omega} := \sum_{n>1} f_n^{G,\omega}, \quad \text{with } f_n^{G,\omega} = \frac{\|f_n\|_{L^2(S^d)}}{\sqrt{\dim(E_n)}} \sum_{\lambda_k^2 = n(n+d-1)} g_{n,k}(\omega) \phi_k,$$

that must be seen as a Gaussian process on the manifold S^d (where $(g_{n,k})_{(n,k)}$ is a sequence of independent standard Gaussian random variables $\mathcal{N}_{\mathbb{R}}(0, 1)$ with $n \in \mathbb{N}^*$ and $\lambda_k^2 = n(n+d-1)$). On a general boundaryless compact Riemannian manifold \mathcal{M} , the Gaussian analogue of (14) should be

$$(21) \quad f^{G,\omega} = \sum_{n>1} f_n^{G,\omega}, \quad \text{with } f_n^{G,\omega} := \frac{\|f_n\|_{L^2(\mathcal{M})}}{\sqrt{\dim(E_{(Kn-K,Kn]})}} \sum_{\lambda_k \in (Kn-K,Kn]} g_{n,k}(\omega) \phi_k.$$

The random functions $f_n^{G,\omega}$ are usually called Gaussian random waves and their study had been introduced by Zelditch in [Zel09]. Let us explain another reason for replacing f_n^ω with $f_n^{G,\omega}$. The rotational invariance of Gaussian vectors implies the following

distribution equivalence (see [Ime19, p. 2731]):

$$(22) \quad f_n^{G,\omega} \sim \frac{1}{\sqrt{\dim(E_{(Kn-K,Kn)})}} \left(\sum_{\lambda_k \in (Kn-K,Kn)} g_{n,k}^2(\omega) \right)^{1/2} f_n^\omega,$$

where all the random variables involved are assumed to be mutually independent. As a consequence of (22), one may directly replace f_n^ω with $f_n^{G,\omega}$ for getting bounds of $\mathbf{E}_\omega[\|f_n^\omega\|_{\mathcal{C}^0(\mathcal{M})}]$ (see Appendix A):

$$\begin{aligned} & \mathbf{E}_\omega[\|f_n^{G,\omega}\|_{\mathcal{C}^0(\mathcal{M})}] \\ &= \frac{1}{\sqrt{\dim(E_{(Kn-K,Kn)})}} \mathbf{E} \left[\left(\sum_{Kn-K < \lambda_k \in Kn} g_{n,k}^2 \right)^{1/2} \right] \times \mathbf{E}_\omega[\|f_n^\omega\|_{\mathcal{C}^0(\mathcal{M})}] \\ &\simeq \mathbf{E}_\omega[\|f_n^\omega\|_{\mathcal{C}^0(\mathcal{M})}]. \end{aligned}$$

Our results will show that the previous equivalence can be generalized to linear combinations of f_n^ω and $f_n^{G,\omega}$ (that is a Riemannian analogue of a result by Marcus and Pisier, see Section 3).

Let us now explain how to deal with (20) or (21). In the classical theory, the main ingredient is the fact that the Gaussian random process (4) is stationary and thus one may use the Dudley-Fernique theorem giving a complete understanding of its almost sure continuity. It seems to be reasonable, at least for the Gaussian random series (20) on spheres S^d , that such ideas can be combined with the transitivity of the isometry group (see an argument at the end of Section 2). It is however clear that such arguments are no longer sufficient for (21) on a general compact Riemannian manifold \mathcal{M} since that case does not seem to fulfill any stationary assumption. However, without a stationary assumption, the Dudley theorem [Dud67] gives a sufficient condition for the almost sure continuity. We refer to the work of Tzvetkov [Tzv10] for a use of the Dudley theorem in the spirit of the Paley-Zygmund theorem (and also [IRT16, Annex 6] for an adaptation of the argument of Tzvetkov).

In order to obtain a necessary and sufficient condition, there are several angles of attack. The first way the author took was to use a generalization of the Dudley-Fernique theorem that weakens the stationary assumption (actually a minor variant of [Dud14, Th. 2.7.4, p. 61]). As pointed out by a referee, one may simplify such a strategy by using the Slepian comparison theorem (see [LT91, Cor. 3.14] or [LQ18b, p. 73]). That is indeed such a proof that will be followed in the sequel: one first completely understands the Gaussian process (20) on the sphere S^d for $d > 2$ (via the Dudley-Fernique theorem) and then the Slepian comparison theorem will show that the almost sure continuity of (20) is equivalent to that of (21) for the manifold setting. In both cases (submanifolds of S^d or of a general manifold), we will obtain sharp estimates of so-called Dudley pseudo-distances whose definition is now recalled.

A fundamental idea of the paper [Dud67] of Dudley can be summarized as follows: the almost sure properties of the Gaussian process $(f^{G,\omega}(x))_{x \in \mathcal{M}}$ are closely related

to the pseudo-distance δ given by

$$(23) \quad \forall (x, y) \in \mathcal{M}^2, \quad \delta(x, y) := \sqrt{\mathbf{E}_\omega[|f^{G,\omega}(x) - f^{G,\omega}(y)|^2]}.$$

Except in very specific examples, one cannot expect to have a simple formula for $\delta(x, y)$. In our concrete situation, the pseudo-distance δ will be equivalent to a simpler distance or pseudo-distance which is a function of the original distance δ_g . An easy computation in (23) indeed shows the formula

$$\delta(x, y)^2 = \sum_{n>1} \|f_n\|_{L^2(\mathcal{M})}^2 \delta_n(x, y)^2,$$

where the partial pseudo-distances δ_n are given by

$$(24) \quad \delta_n(x, y)^2 := \frac{1}{\dim(E_{(Kn-K, Kn]})} \sum_{\lambda_k \in (Kn-K, Kn]} |\phi_k(x) - \phi_k(y)|^2.$$

In another context, a study of those partial pseudo-distances is done by Canzani and Hanin in [CH18] with specific geometric assumptions on the Riemannian manifold \mathcal{M} . In order to dispense with any geometric assumption on \mathcal{M} , we shall combine the following microlocal tools and ideas:

- a part of the work of Canzani and Hanin (in this case Proposition 32 below coming from [CH15b, Lem. 5]) about the derivative of the spectral function on a boundaryless compact manifold,
- choosing K large enough (such an idea is due to Burq and Lebeau in [BL13]) and using off-diagonal estimates obtained by Hörmander (see the proof of Proposition 12 below).

As a consequence, we will prove the equivalence $\delta_n \simeq \min(1, n\delta_g)$ where δ_g is the Riemannian distance of \mathcal{M} (that is a similar form to (8)). Roughly speaking, such an equivalence is possible because concentration of individual eigenfunctions does not matter. The simplicity of such an equivalence is the reason allowing us to simplify the entropy integral in order to recover the Salem-Zygmund condition (19). This article is organized as follows:

- In Section 2, we state the two main theorems concerning the Gaussian random series (20) on S^d and (21) on \mathcal{M} . More precisely, Theorem 2 and Theorem 3 are respectively of qualitative and quantitative nature.
- In Section 3, we first discuss a trivial consequence of Theorem 1 about universality with respect to the choice of X_n in $\sum X_n(\omega) f_n^\omega$ and with respect to the chosen submanifold \mathcal{M}_s . Then we state Theorem 5 ensuring that our analysis is actually independent of the Riemannian metric initially chosen on \mathcal{M} . That result is proved in Section 13 via the semi-classical analysis of the Laplace-Beltrami operator seen as an elliptic differential operator on the compact manifold \mathcal{M} .
- Section 4 is devoted to metric considerations giving a suitable reformulation of the entropy integral.
- In Sections 5 and 6, we prove that the partial pseudo-distances δ_n in (24) are, in some sense, equivalent to the explicit distance $\min(1, n\delta_g)$ (where δ_g stands for

the Riemannian distance). The proofs need to make an accurate comparison with the Bessel function $J_{d/2-1}$. Actually, Theorem 8 and Theorem 10 are the main contributions of the paper that allow to understand the Gaussian processes (20) and (21).

- Section 7 is written for the sake of clarity and contains a few precisions about the Gaussian processes (20) and (21).

- Sections 8, 9 and 10 are devoted to the proofs of the main results in the Gaussian case for spheres and compact manifolds. As written above, we first deal with the sphere case S^d with $d > 2$. Few technical details are necessary because the equivalence $\delta_n \simeq \min(1, n\delta_g)$ does not hold in the whole sphere (for n even, δ_n is not a distance on S^d and does not distinguish antipodal points). The Slepian comparison theorem is then used to deal with submanifolds of S^d or of a general compact manifold \mathcal{M} .

- Sections 11 and 12 contain the proof of Theorem 1 that deals with the initial random series $\sum X_n(\omega)f_n^\omega$. We shall use a truncation argument (already present in the work of Marcus-Pisier) adapted here for the so-called χ distributions.

- Section 13 contains the proof of the invariance with respect to the Riemannian metric of \mathcal{M} (proof of Theorem 5). The proof makes use of the theory of semi-classical pseudo-differential operators.

- Finally, we have gathered two appendices presenting either computations or proofs of more or less known results that we have not found in the literature in the form we need.

Acknowledgements. — The author would like to thank Hervé Que élec for discussions about majorizing measures. Moreover, the author would like to thank the referees who studied the present paper and gave improvements of our initial proofs. Actually, the reference [Dud14, Th. 2.7.4, p. 61], Proposition 6, Proposition 16 and the idea to use the Slepian comparison theorem (see Step 1 in Proposition 17 and the proof in Section 10) are due to them.

2. STATEMENTS OF THE MAIN RESULTS IN THE GAUSSIAN CASE

We now study a necessary and sufficient condition ensuring that the Gaussian random series $\sum f_n^{G,\omega}$ almost surely converges in $\mathcal{C}^0(\mathcal{M})$. It turns out that the condition $\sum_{n>1} \|f_n\|_{L^2(\mathcal{M})}^2 < +\infty$ implies that the Gaussian random series $f^{G,\omega} := \sum_{n>1} f_n^{G,\omega}$ defines an element of $L^2(\Omega)$, hence the Dudley pseudo-distance δ in (23) is well defined on \mathcal{M} (see Proposition 15).

THEOREM 2. — *There is a constant $K_0 > 0$ depending only on the Riemannian manifold \mathcal{M} such that, for any $K > K_0$, if one considers*

- *a sequence of functions $(f_n)_{n>1}$ satisfying $f_n \in E_{(K_n-K, K_n]}$ for each $n \in \mathbb{N}^*$ (see (13)) and $\sum_{n>1} \|f_n\|_{L^2(\mathcal{M})}^2 < +\infty$,*

- *a non-zero dimensional compact submanifold $\mathcal{M}_s \subset \mathcal{M}$ with smooth (eventually empty) boundary,*

then the following assertions are equivalent:

(i) The Gaussian random series $\sum f_n^{G,\omega}$ is almost surely convergent in $\mathcal{C}^0(\mathcal{M}_s)$ (where $f_n^{G,\omega}$ is defined in (21)).

(ii) The Salem-Zygmund condition is fulfilled

$$(25) \quad \sum_{p=1}^{+\infty} \frac{1}{p\sqrt{\ln(p+1)}} \left(\sum_{n=p}^{+\infty} \|f_n\|_{L^2(\mathcal{M})}^2 \right)^{1/2} < +\infty.$$

(iii) Denoting by $\varepsilon \mapsto N_\delta(\mathcal{M}_s, \varepsilon)$ the covering number⁽⁴⁾ function of \mathcal{M}_s with respect to the Dudley pseudo-distance δ defined in (23), then the entropy condition is satisfied

$$(26) \quad \int_0^{+\infty} \sqrt{\ln(N_\delta(\mathcal{M}_s, \varepsilon))} d\varepsilon < +\infty.$$

In the specific case $\mathcal{M} = S^d$, with $d > 2$, the same conclusion holds if each f_n belongs to the eigenspace $\ker(\Delta + n(n+d-1))$ provided that we consider (20) instead of (21).

As for Theorem 1, the almost sure convergence in $\mathcal{C}^0(\mathcal{M})$ is also equivalent to the almost sure convergence in $\mathcal{C}^0(\mathcal{M}_s)$ for any submanifold \mathcal{M}_s .

Let us discuss a quantitative version of the last result. To avoid any problem of measurability due to the uncountability of the submanifold \mathcal{M}_s of \mathcal{M} , one usually sets

$$(27) \quad \mathbf{E}_\omega \left[\sup_{x \in \mathcal{M}_s} \left| \sum_{n>1} f_n^{G,\omega}(x) \right| \right] := \sup_{\substack{\mathcal{F} \subset \mathcal{M}_s \\ \mathcal{F} \text{ countable}}} \mathbf{E}_\omega \left[\sup_{x \in \mathcal{F}} \left| \sum_{n>1} f_n^{G,\omega}(x) \right| \right].$$

In the concrete situation where the four conditions of Theorem 2 are true, the random function $x \mapsto \sum_{n>1} f_n^{G,\omega}(x)$ is almost surely continuous and (27) clearly recovers the classical meaning of $\mathbf{E}_\omega [\| \sum_{n>1} f_n^{G,\omega} \|_{\mathcal{C}^0(\mathcal{M}_s)}]$ by choosing a dense countable subset \mathcal{F} of \mathcal{M}_s . We can now state the quantitative version of Theorem 2.

THEOREM 3. — Under the assumptions of Theorem 2, the expectation

$$\mathbf{E}_\omega \left[\sup_{x \in \mathcal{M}_s} \left| \sum_{n>1} f_n^{G,\omega}(x) \right| \right]$$

and the two numbers appearing in (25) and (26) are equivalent up to a multiplicative loss merely depending on the Riemannian manifold \mathcal{M} , on the submanifold \mathcal{M}_s and on K .

Finally, the conclusion also holds true for the particular case $\mathcal{M} = S^d$ in a similar fashion to the last statement of Theorem 2 (and thus K is irrelevant in the numerical equivalences).

In the specific case $\mathcal{M} = \mathcal{M}_s = S^d$ with $d > 2$, the equivalence of the entropy condition (26) and the expectation (27) can be deduced by considering S^d as a homogeneous space based on the orthogonal group $O_d(\mathbb{R})$. Actually by fixing one point

⁽⁴⁾We recall that $N_\delta(\mathcal{M}_s, \varepsilon)$ is the minimal number of open δ -balls of radius ε that cover \mathcal{M}_s .

$x_0 \in S^d$ then we have

$$\mathbf{E}_\omega \left[\sup_{x \in S^d} \left| \underbrace{\sum_{n>1} f_n^{G,\omega}(x)}_{=f^{G,\omega}(x)} \right| \right] = \mathbf{E}_\omega \left[\sup_{t \in O_d(\mathbb{R})} \left| \underbrace{\sum_{n>1} f_n^{G,\omega}(t \cdot x_0)}_{=F^{G,\omega}(t)} \right| \right]$$

where $F^{G,\omega}$ denotes a Gaussian process on $O_d(\mathbb{R})$ which is stationary (we refer to Proposition 16 for computations showing that $\mathbf{E}[F^{G,\omega}(s)F^{G,\omega}(t)]$ merely depends on $\langle s(x_0), t(x_0) \rangle$ and so merely on $s^{-1}t$). Then the Dudley-Fernique theorem⁽⁵⁾ ensures that the last expectations are equivalent to the entropy condition of $F^{G,\omega}$ on $O_d(\mathbb{R})$. The Dudley pseudo-distances of $(f^{G,\omega}(x))_{x \in S^d}$ and $(F^{G,\omega}(t))_{t \in O_d(\mathbb{R})}$, respectively denoted by δ and δ' , are linked via the formula

$$\forall (s, t) \in O_d(\mathbb{R}) \times O_d(\mathbb{R}), \quad \delta(sx_0, tx_0) = \delta'(s, t).$$

As a consequence, for any $\varepsilon > 0$, we easily see the equality $N_\delta(S^d, \varepsilon) = N_{\delta'}(O_d(\mathbb{R}), \varepsilon)$ of the covering numbers. Hence, the entropy conditions of the Gaussian processes $(f^{G,\omega}(x))_{x \in S^d}$ and $(F^{G,\omega}(t))_{t \in O_d(\mathbb{R})}$ are equivalent.

In order to complete the previous remark, we stress that the main novelty of the paper is that the Dudley pseudo-distance of the a priori non-stationary Gaussian process $(f^{G,\omega}(x))_{x \in \mathcal{M}_s}$ with $\mathcal{M}_s \subset S^d$ or $\mathcal{M}_s \subset \mathcal{M}$, is in some sense equivalent to a much simpler pseudo-distance of the form $\Upsilon(\delta_g)$ (see Theorem 8 and Theorem 10) with $\Upsilon : [0, +\infty) \rightarrow [0, +\infty)$ non-decreasing near 0 and Υ being quite explicit with respect to the coefficients of the random series (see (64)).

3. APPLICATION TO UNIVERSALITY RESULTS FOR RANDOM SERIES

In the statement of Theorem 1, the Salem-Zygmund condition (19) does not involve the random variables X_n provided that the mutual independence and the moment assumption (17) are assumed. Consequently, the almost sure convergence of the random series $\sum X_n(\omega)f_n^\omega(x)$ in $\mathcal{C}^0(\mathcal{M})$ is universal with respect to the random variables (X_n) . That is an analogue of a result by Marcus and Pisier dealing with the random Fourier series $(\omega, x) \in \Omega \times \mathbb{R}/2\pi\mathbb{Z} \mapsto \sum X_n(\omega)e^{inx}$ (see [MP81, p. 7–9] and [Pis78, p. 28, Cor. 7.3]).

For the same reason, Theorem 1 also shows the universality with respect to reasonable submanifolds \mathcal{M}_s of \mathcal{M} although the eigenfunctions are considered with respect to the Laplace-Beltrami operator Δ of the whole Riemannian manifold \mathcal{M} . Indeed with the notations of Theorem 1, we directly see the equivalence of the following two assertions:

- the random series $\sum X_n(\omega)f_n^\omega$ is almost surely convergent in $\mathcal{C}^0(\mathcal{M}_s)$,
- the random series $\sum X_n(\omega)f_n^\omega$ is almost surely convergent in $\mathcal{C}^0(\mathcal{M})$.

⁽⁵⁾Actually we need a non-Abelian Dudley-Fernique theorem that has the same proof as the Abelian case, as noticed in [MP81, p. 96].

For instance, \mathcal{M}_s may be a closed geodesic of \mathcal{M} despite the fact that \mathcal{M}_s is negligible for the Riemannian volume. The previous equivalence contrasts with the classical theory on a compact group. Maybe, the nearest result would be the following one: for any compact Abelian group G and any compact subset $T \subset G$ of *non-empty interior*, then a classical random series almost surely converges in $\mathcal{C}^0(G)$ if and only if it almost surely converges in $\mathcal{C}^0(T)$ (see the argument following [LT91, Th. 13.3]).

We now want to study the universality with respect to the Riemannian metric of \mathcal{M} . We first recall that the Pisier space on \mathbb{T} is the space of functions $\sum_{n \in \mathbb{Z}} c_n e^{inx} \in L^2_x(\mathbb{T})$ such that the associated random Fourier series (4) almost surely converges in $\mathcal{C}^0(\mathbb{T})$ (see [Pis78, p. 2] or [LQ18b, Ch. 6]). In our context, it is clear how to transfer the notion of Pisier space to a boundaryless Riemannian compact manifold \mathcal{M} .

DEFINITION 4. — Given a function $f \in L^2(\mathcal{M})$ and a parameter $K > K_0$, one may decompose

$$f = \left(\int_{\mathcal{M}} f(x) d\mu(x) \right) \mathbf{1}_{\mathcal{M}} + \sum_{n > 1} f_n, \quad \text{with } f_n := \Pi_{(Kn-K, Kn]}(f),$$

where μ is the Riemannian probability measure of \mathcal{M} and $\Pi_{(Kn-K, Kn]} : L^2(\mathcal{M}) \rightarrow L^2(\mathcal{M})$ is the spectral projector on the spectral window $(Kn - K, Kn]$ with respect to $\sqrt{-\Delta}$. Then the “Pisier space on \mathcal{M} ” is the space of functions $f \in L^2(\mathcal{M})$ such that the random series $\sum_{n > 1} f_n^\omega$ almost surely converges in $\mathcal{C}^0(\mathcal{M})$.

For $K \gg 1$, the Salem-Zygmund condition (19) gives an explicit semi-norm on $L^2(\mathcal{M})$ that characterizes the functions $f \in L^2(\mathcal{M})$ of the previous definition. But such a semi-norm clearly involves the spectral decomposition of the Laplace-Beltrami operator Δ which itself is defined via the Riemannian metric of \mathcal{M} . It turns out that the following universality result holds.

THEOREM 5. — *With the above notations, the “Pisier space on \mathcal{M} ” is independent of the spectral parameter K (provided that $K \gg 1$) and of the Riemannian metric of \mathcal{M} .*

Remembering that the space $L^2(\mathcal{M})$ does not depend on the Riemannian structure of \mathcal{M} , one may compare Theorem 5 to the following L^p Paley-Zygmund theorem: for any $p \in [1, +\infty)$, the random series $\sum f_n^\omega$ almost surely converges in $L^p(\mathcal{M})$ if and only if f belongs to $L^2(\mathcal{M})$ (see [Ime19, Th. 2.3] but such a result essentially appears in [BL13] in a different form).

The independence with respect to K in Theorem 5 will be a simple consequence of the Salem-Zygmund condition (19) whereas the independence with respect to the Riemannian metric is more involved and uses

– a semi-classical reformulation of the Salem-Zygmund condition as follows (for a suitable $\Psi \in \mathcal{C}_c^\infty(\mathbb{R})$):

$$\int_0^1 \frac{\|f - \Psi(-h^2 \Delta) f\|_{L^2(\mathcal{M})}}{h \sqrt{-\ln(h)}} dh < +\infty$$

(note that such a reformulation strikingly looks like the condition giving the continuity of stationary Gaussian processes [MS70, Mar73] and Proposition 6),

- a few estimates on semi-classical pseudo-differential operators (see Lemma 27),
- an expansion of the operator $\Psi(-h^2\Delta)$ with respect to h (see the proof of Proposition 30), usually referred as a functional semi-classical calculus, as done by Burq-Gérard-Tzvetkov in [BGT04].

4. METRIC CONSIDERATIONS

In the present section, let us consider a general compact metric space (\mathcal{M}, δ_g) . Although we shall settle general facts, we must have in mind that the next results will be applied in the rest of the paper for \mathcal{M} being a Riemannian compact manifold (or submanifold) with its Riemannian distance δ_g .

PROPOSITION 6. — *Let μ be a Borel probability measure on (\mathcal{M}, δ_g) and let D_g be the diameter of (\mathcal{M}, δ_g) . We moreover assume that there are positive constants H_1, H_2, σ such that for any $t \in [0, D_g]$ and $x \in \mathcal{M}$ we have*

$$(28) \quad H_1 \left(\frac{t}{D_g} \right)^\sigma \subset \mu(B_{\delta_g}(x, t)) \subset H_2 \left(\frac{t}{D_g} \right)^\sigma.$$

Let $\Upsilon : [0, +\infty) \rightarrow [0, +\infty)$ be a subadditive, non-decreasing and right-continuous function satisfying $\Upsilon(0) = 0$ and a pseudo-distance δ on \mathcal{M} which is equivalent to the pseudo-distance $\Upsilon(\delta_g)$ as follows:

$$(29) \quad \exists \varrho > 1, \quad \frac{\delta}{\sqrt{\varrho}} \subset \Upsilon(\delta_g) \subset \sqrt{\varrho} \delta.$$

Denoting by $N_\delta(\mathcal{M}, \varepsilon)$ the minimal number of δ -open balls of radius ε that cover \mathcal{M} , then the entropy condition for δ

$$(30) \quad \int_0^{+\infty} \sqrt{\ln(N_\delta(\mathcal{M}, \varepsilon))} d\varepsilon < +\infty$$

is equivalent to

$$\int_0^1 \frac{\Upsilon(t)}{t\sqrt{\ln(1/t)}} dt < +\infty.$$

Proof

Topological remark. — One may assume that Υ does not identically vanish, otherwise the conclusion is obvious and of no interest. We also note that the subadditivity assumption ensures that $\Upsilon(t) > 0$ for any $t > 0$ and then $\Upsilon(\delta_g)$ turns out to be a distance. We also easily infer that δ and δ_g give rise to the same topology on \mathcal{M} .

Step 1. — We now introduce a pseudo-inverse function $\Upsilon^{-1} : [0, +\infty) \rightarrow [0, +\infty]$ as follows

$$\forall \varepsilon > 0, \quad \Upsilon^{-1}(\varepsilon) := \inf\{t > 0, \Upsilon(t) > \varepsilon\},$$

with the usual convention $\inf \emptyset = +\infty$. The right-continuity of Υ implies the following two equivalences

$$\begin{aligned}\Upsilon(t) > \varepsilon &\iff t > \Upsilon^{-1}(\varepsilon), \\ \Upsilon(t) < \varepsilon &\iff t < \Upsilon^{-1}(\varepsilon).\end{aligned}$$

Hence, we are able to come back to the open balls for the original distance δ_g . For any $x \in \mathcal{M}$ and any $\varepsilon > 0$ we have

$$\begin{aligned}B_{\Upsilon(\delta_g)}(x, \varepsilon) &= B_{\delta_g}(x, \Upsilon^{-1}(\varepsilon)), \\ \mu(B_{\Upsilon(\delta_g)}(x, \varepsilon)) &= \mu(B_{\delta_g}(x, \Upsilon^{-1}(\varepsilon))).\end{aligned}$$

Let us set $H = H_2/H_1$. The last equality, (28) and (29) easily imply that μ is also almost-homogeneous for δ in the following sense (in which we note the multiplicative loss ϱ in the right-hand side):

$$(31) \quad \forall (x, y) \in \mathcal{M}^2, \forall \varepsilon > 0, \quad \mu(B_\delta(x, \varepsilon)) \triangleleft H\mu(B_\delta(y, \varrho\varepsilon)).$$

Step 2. — The following consequence of Step 1 is well-known: for any $\varepsilon > 0$ and any $x \in \mathcal{M}$ we have $\mu(B_\delta(x, \varepsilon)) > 0$ and

$$(32) \quad \frac{1}{H\mu(B_\delta(x, \varrho\varepsilon))} \triangleleft N_\delta(\mathcal{M}, \varepsilon) \triangleleft \frac{H}{\mu(B_\delta(x, \varepsilon/2\varrho))}.$$

For the bound from below, we consider x_1, \dots, x_N with $N = N_\delta(\mathcal{M}, \varepsilon)$ so that the open ball $B_\delta(x, \varepsilon)$ cover \mathcal{M} . By the subadditivity of the probability measure μ (from the previous topological remark, $B_\delta(x, \varepsilon)$ is also open for the original topology on \mathcal{M}), (31) and the equality

$$\mathcal{M} := B_\delta(x_1, \varepsilon) \cup \dots \cup B_\delta(x_N, \varepsilon)$$

we get the bound from below of (32). For the bound from above in (32), let N^* be the so-called packing number defined as the largest integer such that (\mathcal{M}, δ) contains N^* disjoint open balls $B_\delta(x_1, \varepsilon/2), \dots, B_\delta(x_{N^*}, \varepsilon/2)$. It is well known that $N_\delta(\mathcal{M}, \varepsilon) \triangleleft N^*$. We then easily obtain (32) thanks to (31) and the inequalities

$$\frac{1}{H} N_\delta(\mathcal{M}, \varepsilon) \mu(B_\delta(x, \varepsilon/2\varrho)) \triangleleft \sum_{k=1}^{N^*} \mu(B_\delta(x_k, \varepsilon/2)) \triangleleft 1.$$

Step 3. — Since D_g is diameter of (\mathcal{M}, δ_g) , we have $\delta(x) \triangleleft \sqrt{\varrho} \Upsilon(D_g)$ for any $x \in \mathcal{M}$, and so for any $\varepsilon > \sqrt{\varrho} \Upsilon(D_g)$ we have $N_\delta(\mathcal{M}, \varepsilon) = 1$. That fact leads to the well-known remark stating that the entropy condition (30) is merely relevant for $\varepsilon \rightarrow 0$.

In other words, for a fixed element $x \in \mathcal{M}$, the inequalities (32) show that the entropy condition (30) is equivalent to the finiteness of

$$\int_0^{+\infty} \sqrt{\ln \left(\frac{1}{\mu(B_{\delta_g}(x, \Upsilon^{-1}(\varepsilon)))} \right)} d\varepsilon = \int_0^{+\infty} \sqrt{\ln \left(\frac{1}{\mu(B_\delta(x, \varepsilon))} \right)} d\varepsilon.$$

Note that the two last integrals are indeed supported in $[0, \Upsilon(D_g)]$ and their finiteness is also relevant at $\varepsilon \rightarrow 0$. By using (28), their finiteness is then equivalent to the finiteness of the following integrals (the second one is in the sense of Stieltjes):

$$(33) \quad \int_0^{\Upsilon(D_g)} \sqrt{\ln(D_g/\Upsilon^{-1}(\varepsilon))} d\varepsilon = \int_{(0, D_g]} \sqrt{\ln(D_g/t)} d\Upsilon(t).$$

For any $a \in (0, D_g)$, an integration by parts in the sense of Stieltjes and the condition $\Upsilon(0) = 0$ give

$$\frac{1}{2} \int_a^{D_g} \frac{\Upsilon(t)}{\sqrt{\ln(D_g/t)}} \frac{dt}{t} = \sqrt{\ln(D_g/a)} \underbrace{\Upsilon(a)}_{=\int_{(0, a]} d\Upsilon(t)} + \int_{(a, D_g]} \sqrt{\ln(D_g/t)} d\Upsilon(t).$$

By using the last computations and the inequality $\sqrt{\ln(D_g/a)} \ll \sqrt{\ln(D_g/t)}$ for any $t \in (0, a]$, we obtain

$$\int_{(a, D_g]} \sqrt{\ln(D_g/t)} d\Upsilon(t) \ll \frac{1}{2} \int_a^{D_g} \frac{\Upsilon(t)}{\sqrt{\ln(D_g/t)}} \frac{dt}{t} \ll \int_{(0, D_g]} \sqrt{\ln(D_g/t)} d\Upsilon(t).$$

By making a tend to 0^+ and looking at (33), we get

$$\int_0^{\Upsilon(D_g)} \sqrt{\ln(D_g/\Upsilon^{-1}(\varepsilon))} d\varepsilon = \frac{1}{2} \int_0^1 \frac{\Upsilon(D_g t)}{t \sqrt{\ln(1/t)}} dt.$$

Step 4. — We now explain how to get rid of the parameter D_g in the last integral. For any $s > 0$, we denote by $\lceil s \rceil$ the least integer greater than or equal to a given number $s > 0$. By using the fact that Υ is subadditive and non-decreasing, one remarks the inequality $\Upsilon(st) \ll \lceil s \rceil \Upsilon(t)$ for any $t > 0$. By replacing (s, t) with $(1/s, st)$, the previous bound can be reversed as follows

$$(34) \quad \frac{1}{\lceil 1/s \rceil} \Upsilon(t) \ll \Upsilon(st) \ll \lceil s \rceil \Upsilon(t).$$

In other words, one may replace $\Upsilon(D_g t)$ with $\Upsilon(t)$ provided we authorize a multiplicative loss merely depending on D_g . The proof is finished.

Note that the above proof indeed shows the quantitative following version:

COROLLARY 7. — *Under the assumptions of Proposition 6, we have the equivalence*

$$\int_0^{+\infty} \sqrt{\ln(N_\delta(\mathcal{M}, \varepsilon))} d\varepsilon \simeq \int_0^1 \frac{\Upsilon(t)}{t \sqrt{\ln(1/t)}} dt$$

up to multiplicative constants which are independent of Υ (but possibly dependent on $D_g, H_2/H_1$ and ϱ).

Proof. — By looking at the proof of Proposition 6, we have to check in Step 3 that $\Upsilon(D_g)$ is controlled by any of the two integrals with a constant which is independent of Υ . For the entropy integral, let us denote by D the diameter of (\mathcal{M}, δ) . We have $\Upsilon(D_g) \ll \sqrt{\varrho} D$ thanks to (29). Since no ball of radius $D/3$ can cover (\mathcal{M}, δ) , we get

$$\int_0^{+\infty} \sqrt{\ln(N(\mathcal{M}, \varepsilon))} d\varepsilon > \int_0^{D/3} \sqrt{\ln(N(\mathcal{M}, \varepsilon))} d\varepsilon > \sqrt{\ln(2)} \frac{D}{3} > \frac{\sqrt{\ln(2)}}{3\varrho} \Upsilon(D_g).$$

For the integral $\int_0^1 (\Upsilon(t)/t\sqrt{\ln(1/t)}) dt$, we use (34) as follows:

$$\int_{1/2}^1 \frac{\Upsilon(t)}{t\sqrt{\ln(1/t)}} dt > 2\sqrt{\ln(2)} \Upsilon(1/2) > \frac{2\sqrt{\ln(2)}}{[2D_g]} \Upsilon(D_g).$$

5. PARTIAL PSEUDO-DISTANCES ON SPHERES

For any integers $d > 2$ and $n > 1$, we denote by δ_n the pseudo-distance on the sphere S^d given by

$$(35) \quad \delta_n(x, y)^2 := \frac{1}{\dim(E_n)} \sum_{\lambda_k^2 = n(n+d-1)} |\phi_k(x) - \phi_k(y)|^2,$$

for any $(x, y) \in S^d \times S^d$ and where E_n stands for the eigenspace $\ker(\Delta + n(n+d-1))$. It is well-known that E_n is the space of the restrictions to S^d of the n -homogeneous harmonic polynomials of \mathbb{R}^{d+1} (see for instance [Far08, Prop. 9.3.5]). In particular, for n even, the relation $\phi_k(x) = \phi_k(-x)$ holds for any $\phi_k \in E_n$ and $x \in S^d$. And thus we get $\delta_n(x, -x) = 0$ for n even. In other words, if one wants to compare δ_n to the Riemannian distance δ_g of S^d , one must unavoidably restrict δ_n to a reasonable subset of S^d that does not contain antipodal points. These considerations lead us to the following statement (in which $\overline{B_{\delta_g}}(p, \vartheta)$ is a Riemannian closed ball of S^d).

THEOREM 8. — *For any $d > 2$ and for any $\vartheta \in (0, \frac{\pi}{2})$, there is $C = C(d, \vartheta) > 1$ such that for any $n \in \mathbb{N}^*$ and any $p \in S^d$, the pseudo-distance δ_n is equivalent on the closed ball $\overline{B_{\delta_g}}(p, \vartheta)$ to the distance $\min(1, n\delta_g)$ as follows*

$$\frac{1}{C} \min(1, n\delta_g) \circlearrowleft \delta_n \circlearrowright C \min(1, n\delta_g).$$

Proof. — Let us denote by $e_n(x, y)$ the reproducing kernel of the eigenspace $\ker(\Delta + n(n+d-1))$:

$$e_n(x, y) := \sum_{\lambda_k^2 = n(n+d-1)} \phi_k(x)\phi_k(y),$$

where each eigenfunction ϕ_k is assumed to be real-valued. The pseudo-distance δ_n then takes the form

$$(36) \quad \delta_n(x, y)^2 = \frac{e_n(x, x) + e_n(y, y) - 2e_n(x, y)}{\dim(E_n)}.$$

Let us recall how $e_n(x, y)$ may be expressed thanks to orthogonal polynomials (for instance Gegenbauer polynomials). We prefer here Jacobi polynomials (which are directly related to Gegenbauer polynomials). Let $(P_n^{(d/2-1, d/2-1)})_{n \in \mathbb{N}}$ be the family of Jacobi polynomials associated to the weight $t \in [-1, 1] \mapsto (1-t^2)^{d/2-1}$ and satisfying (see [Sze75, p. 58]):

$$(37) \quad P_n^{(d/2-1, d/2-1)}(1) = \binom{n+d/2-1}{n} \underset{n \rightarrow +\infty}{\sim} \frac{n^{d/2-1}}{\Gamma(d/2)}.$$

The so-called additional formula (see [SW71, Lem. 2.8 (p. 143) & Th. 2.14 (p. 149)]) ensures the existence of a constant $c_{d,n}$ satisfying

$$(38) \quad e_n(x, y) = c_{d,n} P_n^{(d/2-1, d/2-1)}(\langle x, y \rangle).$$

Note that $e_n(x, x) = c_{d,n} P_n^{(d/2-1, d/2-1)}(1)$ does not depend on x and so equals

$$\int_{S^d} e_n(x, x) \frac{dx}{\text{vol}_{S^d}(S^d)} = \frac{\dim(E_n)}{\text{vol}_{S^d}(S^d)}.$$

As a consequence, one has

$$(39) \quad \frac{c_{d,n}}{\dim(E_n)} = \frac{1}{\text{vol}_{S^d}(S^d) P_n^{(d/2-1, d/2-1)}(1)} \simeq \frac{1}{n^{d/2-1}}.$$

We now simplify (36) as $(2/\dim(E_n))(e_n(x, x) - e_n(x, y))$ and we get the following closed form

$$(40) \quad \delta_n(x, y)^2 = \frac{2c_{d,n}}{\dim(E_n)} \left[P_n^{(d/2-1, d/2-1)}(1) - P_n^{(d/2-1, d/2-1)}(\langle x, y \rangle) \right].$$

One may write $\langle x, y \rangle = \cos(\delta_g(x, y))$ where $\delta_g(x, y) \in [0, \pi]$ is the Riemannian distance between x and y on the sphere S^d . Hence, we have the equivalence

$$(41) \quad 1 - \langle x, y \rangle \simeq \delta_g(x, y)^2.$$

Note that the Riemannian distance between two elements x and y on the spherical cap $\overline{B_{\delta_g}(p, \vartheta)}$ is less or equal to 2ϑ . Hence we have

$$\langle x, y \rangle = \cos(\delta_g(x, y)) > \cos(2\vartheta) > -1.$$

So the conclusion is a consequence of Proposition 9 below and of (39), (40) and (41).

PROPOSITION 9. — *For any integer $d > 2$ and any real number $\vartheta \in (0, \pi/2)$, there is a constant $C_{d,\vartheta} > 1$ such that the following inequalities hold true for any $n > 1$ and any $\alpha \in [\cos(2\vartheta), 1)$*

$$(42) \quad \frac{1}{C_{d,\vartheta}} \circlearrowleft \frac{P_n^{(d/2-1, d/2-1)}(1) - P_n^{(d/2-1, d/2-1)}(\alpha)}{n^{d/2-1} \min(1, n^2(1 - \alpha))} \circlearrowright C_{d,\vartheta}.$$

Proof

Step 1. — We first prove a weak version of (42) in which the constant $C_{d,\vartheta}$ may depend on n . We set

$$Q_{n,d}(\alpha) := \frac{P_n^{(d/2-1, d/2-1)}(1) - P_n^{(d/2-1, d/2-1)}(\alpha)}{1 - \alpha},$$

which is a polynomial with respect to α . It turns out that $P_n^{(d/2-1, d/2-1)}$ reaches its maximum in $[\cos(2\vartheta), 1]$ at the mere point 1 (see [Sze75, p. 168]) and so the polynomial $Q_{n,d}$ is positive on $[\cos(2\vartheta), 1)$. Moreover, it also does not vanish for $\alpha = 1$ thanks to the following formula (see (37) and [Sze75, p. 63, (4.21.7)]):

$$(43) \quad (P_n^{(d/2-1, d/2-1)})'(1) = \frac{n + d - 1}{2} P_{n-1}^{(d/2, d/2)}(1) > 0.$$

By compactness of $[\cos(2\vartheta), 1]$, there is a constant $C_{d,\vartheta,n} > 1$ and such that the following holds for any $\alpha \in [\cos(2\vartheta), 1]$

$$\frac{1}{C_{d,\vartheta,n}} \circlearrowleft Q_{n,d}(\alpha) \circlearrowright C_{d,\vartheta,n}.$$

Since $1 - \alpha$ belongs to $[0, 2]$, we have

$$\frac{\min(1, n^2(1 - \alpha))}{n^2} \oslash (1 - \alpha) \oslash 2 \min(1, n^2(1 - \alpha))$$

and hence

$$\frac{1}{n^2 C_{d, \vartheta, n}} \oslash \frac{P_n^{(d/2-1, d/2-1)}(1) - P_n^{(d/2-1, d/2-1)}(\alpha)}{\min(1, n^2(1 - \alpha))} \oslash 2 C_{d, \vartheta, n}.$$

Such estimates are very far from the expected inequalities but show that it is sufficient to prove (42) for $n > n_{d, \vartheta}$ (for a suitable positive integer $n_{d, \vartheta}$ merely depending on (d, ϑ)).

Step 2. — The mean value theorem and the formula giving the derivative of a Jacobi polynomial (as in (43)) ensure the existence of $z \in (\alpha, 1)$ such that the previous computations can be continued:

$$Q_{n,d}(\alpha) = \frac{d}{dz} P_n^{(d/2-1, d/2-1)}(z) = \frac{n+d-1}{2} P_{n-1}^{(d/2, d/2)}(z).$$

Classical estimates on Jacobi polynomials (see the proof of [Ime18, Lem. 3.9] or (47) below) ensure that there is $c_1 > 0$, merely depending on the dimension d , such that the equivalence $P_{n-1}^{(d/2, d/2)}(\cos(\theta)) \simeq n^{d/2}$ uniformly holds true with respect to $\theta \in [0, c_1/n]$. As a consequence, we get

$$\forall n \gg 1, \forall \theta \in (0, c_1/n], \quad \frac{Q_{d,n}(\cos(\theta))}{n^2} \simeq n^{d/2-1}.$$

In the last regime, we remark the equivalence $n^2(1 - \cos(\theta)) \simeq \min(1, n^2(1 - \cos(\theta)))$. We thus have proved

$$(44) \quad \forall n \gg 1, \forall \theta \in (0, c_1/n], \quad \frac{P_n^{(d/2-1, d/2-1)}(1) - P_n^{(d/2-1, d/2-1)}(\cos(\theta))}{\min(1, n^2(1 - \cos(\theta)))} \simeq n^{d/2-1}.$$

Step 3. — Thanks to [Sze75, Th. 7.32.2, p. 169], we know that for any constant $c_2 > 0$ one may find $C_2 > 1$ such that

$$\forall n \gg 1, \forall \theta \in [c_2/n, \pi/2], \quad |P_n^{(d/2-1, d/2-1)}(\cos(\theta))| \oslash \frac{C_2}{\sqrt{n} \theta^{(d-1)/2}}.$$

The formula $P_n^{(d/2-1, d/2-1)}(-x) = (-1)^n P_n^{(d/2-1, d/2-1)}(x)$ (see [Sze75, p. 59]) and the previous inequality implies the following one for any constant c_3 larger than c_2 :

$$\forall n \gg 1, \forall \theta \in [c_3/n, \pi - c_3/n], \quad |P_n^{(d/2-1, d/2-1)}(\cos(\theta))| \oslash C_2 \frac{n^{d/2-1}}{c_3^{(d-1)/2}}.$$

Remembering (37) and noting that C_2 is independent of c_3 , one sees that one may choose c_3 large enough satisfying for all $\theta \in [c_3/n, \pi - c_3/n]$:

$$(45) \quad P_n^{(d/2-1, d/2-1)}(1) - P_n^{(d/2-1, d/2-1)}(\cos(\theta)) \underset{n \rightarrow +\infty}{\simeq} n^{d/2-1}.$$

Since we assumed $\vartheta < \pi/2$, the inequality $2\vartheta \in \pi - c_3/n$ holds true for $n > n_{d,\vartheta}$ (with a suitable positive integer $n_{d,\vartheta}$). In contrast with (44), we have the equivalence $\min(1, n^2(1 - \cos(\theta))) \simeq 1$ in such a regime. As a consequence, we have

$$(46) \quad \forall n > n_{d,\vartheta}, \forall \theta \in [c_3/n, 2\vartheta],$$

$$\frac{P_n^{(d/2-1, d/2-1)}(1) - P_n^{(d/2-1, d/2-1)}(\cos(\theta))}{\min(1, n^2(1 - \cos(\theta)))} \simeq n^{d/2-1}.$$

Step 4. — Comparing (44) and (46), one cannot exclude the possibility that c_3 may have been chosen too large in Step 3. To complete our proof, we need to understand the case $c_1 < c_3$. Here we invoke the following uniform limit with respect to $t \in [0, c_3]$ (see [Sze75, p. 192, (8.1.1)]) that makes a connexion between the Jacobi polynomials $P_n^{(d/2-1, d/2-1)}$ and the Bessel function $J_{d/2-1}$:

$$(47) \quad \lim_{n \rightarrow +\infty} \frac{P_n^{(d/2-1, d/2-1)}(\cos(t/n))}{n^{d/2-1}} = 2^{d/2-1} \frac{J_{d/2-1}(t)}{t^{d/2-1}}$$

in the Banach space $\mathcal{C}_t^0([0, c_3], \mathbb{R})$.

The Poisson representation of the Bessel function $J_{d/2-1}$ (see [Sze75, p. 15, (1.71.6)]) suggests introducing the following real-valued function

$$(48) \quad \mathfrak{F}_{d-1}(t) := \frac{J_{d/2-1}(t)}{t^{d/2-1}} = \frac{1}{2^{d/2-1} \Gamma((d-1)/2) \sqrt{\pi}} \int_{-1}^1 e^{its} (1-s^2)^{(d-3)/2} ds.$$

The notation \mathfrak{F}_{d-1} is justified since that function may be interpreted as the radial part of the Fourier transform of the spherical measure σ_{d-1} on S^{d-1} (see (56)). One directly checks that the function \mathfrak{F}_{d-1} is real-valued, admits a maximum at the unique point $t = 0$ and satisfies $\lim_{t \rightarrow +\infty} \widehat{\sigma}_{d-1}(t) = 0$ (thanks to the Riemann-Lebesgue lemma). As a consequence, we get the strict inequality $\sup_{t > c_1} \widehat{\sigma}_{d-1}(t) < \widehat{\sigma}_{d-1}(0)$ which in turn implies the equivalent one

$$(49) \quad \exists \rho \in (0, 1), \forall t > c_1, \widehat{\sigma}_{d-1}(t) < \rho \widehat{\sigma}_{d-1}(0).$$

Coming back to (47), we see that the sequence of functions

$$t \mapsto \frac{1}{n^{d/2-1}} \left[P_n^{(d/2-1, d/2-1)}(\cos(t/n)) - \rho P_n^{(d/2-1, d/2-1)}(1) \right]$$

uniformly converges on $[c_1, c_3]$ to the continuous negative function

$$t \mapsto 2^{d/2-1} [\widehat{\sigma}_{d-1}(t) - \rho \widehat{\sigma}_{d-1}(0)].$$

Hence, we infer that the following holds true

$$\forall n \gg 1, \forall \theta \in [c_1/n, c_3/n], P_n^{(d/2-1, d/2-1)}(\cos(\theta)) \in \rho P_n^{(d/2-1, d/2-1)}(1).$$

Such new estimates can now be combined to

$$-P_n^{(d/2-1, d/2-1)}(\cos(\theta)) \in C_d P_n^{(d/2-1, d/2-1)}(1)$$

(as a consequence of (47)). Hence (45) and (46) hold with c_1 (instead of c_3). Taking account of (44), we finally obtain the wanted uniform equivalence (42).

6. PARTIAL PSEUDO-DISTANCES ON MANIFOLDS

In this part, \mathcal{M} is a general boundaryless compact Riemannian manifold (and we use the notations of the introduction about the spectral analysis of the Laplace-Beltrami operator). For any interval $\mathcal{J} \subset [0, +\infty)$ we define the spectral function of \mathcal{J} (with respect to $\sqrt{-\Delta}$) as follows: for any $(x, y) \in \mathcal{M}^2$ we set

$$e_{\mathcal{J}}(x, y) = \sum_{\lambda_k \in \mathcal{J}} \phi_k(x) \phi_k(y),$$

where we recall that each eigenfunction ϕ_k of Δ is assumed to be real-valued. Let us recall an idea we attribute to Burq and Lebeau in [BL13, p. 923] (we also refer to the end of the proof of [Ime19, Lem. 8.1] for details): one may find two numbers $K_0 > 1$ and $C > 1$ that merely depend on the Riemannian manifold \mathcal{M} such that the following on-diagonal estimates hold

$$(50) \quad \forall K > K_0, \forall n > 1, \forall x \in \mathcal{M}, \quad \frac{K^d n^{d-1}}{C} \ll e_{(Kn-K, Kn]}(x, x) \ll CK^d n^{d-1}.$$

Upon modifying C , we note that the same estimates hold for $\dim(E_{(Kn-K, Kn]})$ by integration on the compact manifold \mathcal{M} :

$$(51) \quad \forall K > K_0, \forall n > 1, \forall x \in \mathcal{M}, \quad \frac{K^d n^{d-1}}{C} \ll \dim(E_{(Kn-K, Kn]}) \ll CK^d n^{d-1}.$$

We now introduce a pseudo-distance δ_n on the manifold \mathcal{M} which is analogue to (35) (we keep the same notation δ_n for simplicity). For any pair $(x, y) \in \mathcal{M}^2$ we set

$$(52) \quad \delta_n(x, y)^2 := \frac{1}{\dim(E_{(Kn-K, Kn]})} \sum_{\lambda_k \in (Kn-K, Kn]} |\phi_k(x) - \phi_k(y)|^2$$

$$(53) \quad = \frac{e_{(Kn-K, Kn]}(x, x) + e_{(Kn-K, Kn]}(y, y) - 2e_{(Kn-K, Kn]}(x, y)}{\dim(E_{(Kn-K, Kn]})}.$$

By bounding $|\phi_k(x) - \phi_k(y)|^2 \ll 2(|\phi_k(x)|^2 + |\phi_k(y)|^2)$ and using (50) and (51), we remark the uniform estimate

$$(54) \quad \sup_{n > 1} \sup_{(x, y) \in \mathcal{M}^2} \delta_n(x, y) < +\infty.$$

In the sequel, we will prove the following analogue result of Theorem 8 (but here the equivalence (55) holds in the whole manifold \mathcal{M}).

THEOREM 10. — *There are two constants $K_0 > 1$ and $C > 1$ that merely depend on the Riemannian manifold \mathcal{M} such that the following equivalence⁽⁶⁾ holds true for any $K > K_0$, any pair $(x, y) \in \mathcal{M}^2$ and any $n \in \mathbb{N}^*$:*

$$(55) \quad \frac{1}{C} \min(1, Kn\delta_g(x, y)) \ll \delta_n(x, y) \ll C \min(1, Kn\delta_g(x, y)).$$

⁽⁶⁾We note that $\delta_n(x, y)$ implicitly depends on K .

The main interest of the previous result is that it holds true without any geometric assumption on the boundaryless compact Riemannian manifold \mathcal{M} . It indeed relies on the freedom to set K large enough. Under specific geometric assumptions, a similar upper bound to that of (55) appears in [CH15a, Th. 1]. Moreover, a much more precise than (55) is given in [CH15b, Th. 9] by assuming a “geometric mutually nonfocal hypothesis” on the manifold \mathcal{M} .

We now recall the relation linking the Bessel function $J_{d/2-1}$ and the Fourier transform of the spherical measure σ_{d-1} on S^{d-1} , for any $\nu \in \mathbb{R}^d \setminus \{0\}$ we set

$$\frac{J_{(d-2)/2}(|\nu|)}{|\nu|^{(d-2)/2}} = \frac{1}{(2\pi)^{d/2}} \int_{S^{d-1}} e^{i\langle \nu, w \rangle} d\sigma_{d-1}(w) = \frac{1}{(2\pi)^{d/2}} \int_{S^{d-1}} \exp(i|\nu|w_1) d\sigma_{d-1}(w).$$

Since the previous Fourier transform is radial, we make the following abuse of notation:

$$(56) \quad \forall \nu > 0, \quad \mathfrak{F}_{d-1}(\nu) := \frac{J_{(d-2)/2}(\nu)}{\nu^{(d-2)/2}} = \frac{1}{(2\pi)^{d/2}} \int_{S^{d-1}} \exp(i\nu w_1) d\sigma_{d-1}(w).$$

We will use some material developed by Canzani and Hanin. More precisely we need a suitable asymptotic of the spectral function given in the next theorem (proved by Hörmander in [Hör68, Th. 4.4] for the case $|I| = |J| = 0$). But the choice of the explicit principal term has been enlightened by Canzani and Hanin in [CH15b]. The asymptotic formula, involving derivatives, is now considered as known (see the introduction of [CH18]). We give here a reformulation with coordinate patches.

THEOREM 11. — *There is $\alpha_0 > 0$ such that for any coordinate patch $\tau : U \subset \mathbb{R}^d \rightarrow V \subset \mathcal{M}$ with $\text{diam}(V) < \alpha_0$, for any multi-indexes $I \in \mathbb{N}^d$ and $J \in \mathbb{N}^d$, the following asymptotic holds true for any $(x, y) \in V^2$ and $\lambda > 0$ (where the derivatives are seen in the coordinate patch):*

$$\partial_x^I \partial_y^J \sum_{\lambda_j \in \lambda} \phi_j(x) \phi_j(y) = \frac{\partial_x^I \partial_y^J}{(2\pi)^{d/2}} \int_0^\lambda \nu^{d-1} \mathfrak{F}_{d-1}(\nu \delta_g(x, y)) d\nu + \mathcal{O}((1 + \lambda)^{d-1+|I|+|J|}).$$

Finally, the remainder is uniform provided that x and y run over a compact subset of the open set V .

Proof. — Since we do not know a published reference where Theorem 11 is stated with a proof, we explain in Appendix B how it can be easily recovered as a consequence of the analysis of [CH18], a Bernstein-type inequality proved in [Bin04] and a Tauberian theorem as in [Sog17, Ch. 4].

The reason of restricting the estimates of Theorem 11 to compact subsets of the open subset V is due to the definition of the $\mathcal{C}^\infty(\mathcal{M})$ -topology (see [Die72, Ch. XVII.1 & XVII.2]). Obviously for $|I| = |J| = 0$, coordinate patches are useless and the estimates hold on the whole compact manifold.

If the two points x and y are far from each other, we shall need to forget the principal term of the asymptotic of the spectral function (at least for $|I| = |J| = 0$). In [CH15b, line (79)], a geometric mutually nonfocal hypothesis on the manifold \mathcal{M} is assumed, and thus Canzani and Hanin make use of an asymptotic of Safarov

[Saf88] stating $e_{[0,\lambda]}(x, y) = o(\lambda^{d-1})$. Since our issue is posed without any geometric assumption, we replace their argument with the o -diagonal Hörmander estimate $e_{[0,\lambda]}(x, y) = \mathcal{O}(\lambda^{d-1})$.

PROPOSITION 12. — *For any fixed $\alpha > 0$, one may find two numbers $K_0 > 1$ and $c > 1$ (depending on the Riemannian manifold \mathcal{M} and $\alpha > 0$) satisfying the following property: for any $K > K_0$ the following implication holds true for any $n \in \mathbb{N}^*$ and any pair $(x, y) \in \mathcal{M}^2$:*

$$\delta_g(x, y) > \alpha \implies \frac{1}{c} \delta_n(x, y) \delta c.$$

Proof. — The upper bound $\delta_n(x, y) \delta c$ has already been proved in (54). From [Hör68, line (4.11)] (see also [Shu01, p. 162, Th. 21.1, point 2]), we know that for any $\alpha > 0$ there is $C > 0$ such that, for any $(x, y) \in \mathcal{M}^2$, the following uniform o -diagonal estimate holds true

$$\delta_g(x, y) > \alpha \implies \forall \lambda > 0, |e_{[0,\lambda]}(x, y)| \delta C(1 + \lambda)^{d-1},$$

where C depends on α and on the Riemannian structure of \mathcal{M} . Thus we obtain

$$\begin{aligned} |e_{(K_n-K, K_n)}(x, y)| &= |e_{[0, K_n+K]}(x, y) - e_{[0, K_n]}(x, y)| \\ &\delta C(1 + K_n + K)^{d-1} + C(1 + K_n)^{d-1} \\ &\delta C(3^{d-1} + 2^{d-1})(K_n)^{d-1}. \end{aligned}$$

We now recall the estimate $e_{(K_n-K, K_n)}(x, x) \simeq K^d n^{d-1}$ (see (50)), which also implies $\dim(E_{(K_n-K, K_n)}) \simeq K^d n^{d-1}$ by integration over \mathcal{M} . Remembering the formula (53), we infer that there are three positive constants C_1, C_2, C_3 (independent of n and K) such that

$$(57) \quad \begin{aligned} \delta_n(x, y)^2 &> \frac{n^{d-1}}{\dim(E_{(K_n-K, K_n)})} [C_1 K^d - C_2 K^{d-1}] \\ &> \frac{C_1 K^d - C_2 K^{d-1}}{C_3 K^d}. \end{aligned}$$

We conclude by making K tend to $+\infty$.

We now need to improve the previous result if $n\delta_g(x, y)$ is bounded from below or from above. By comparison with [CH15b, p. 1728–1729] that involves geometric assumptions on the manifold \mathcal{M} and relies on several tools obtained by Zelditch, Potash and Xu, our proofs of the next two results merely use the asymptotic of the spectral function given by Theorem 11.

PROPOSITION 13. — *For any fixed $\beta > 0$, one may find two numbers $K_0 > 1$ and $c > 1$ (depending on the Riemannian manifold \mathcal{M} and $\beta > 0$) satisfying the following property: for any $K > K_0$, for any $n \in \mathbb{N}^*$ and any pair $(x, y) \in \mathcal{M}^2$ the following implication holds true*

$$\delta_g(x, y) > \frac{\beta}{Kn} \implies \frac{1}{c} \delta_n(x, y) \delta c.$$

Proof. — As for Proposition 12, the upper bound $\delta_n \in c$ is given in (54). To get the lower bound, we apply Proposition 12 with the constant $\alpha = \alpha_0$ appearing in the statement of Theorem 11. Hence, the case $\delta_g(x, y) > \alpha_0$ is already done. We now assume $\alpha_0 > \delta_g(x, y) > \beta/Kn$ and we have the following asymptotic

$$(58) \quad e_{(Kn-K, Kn]}(x, y) = \int_{Kn-K}^{Kn} \nu^{d-1} \mathfrak{F}_{d-1}(\nu \delta_g(x, y)) \frac{d\nu}{(2\pi)^{d/2}} + (Kn)^{d-1} \mathcal{O}(1),$$

where the remainder $\mathcal{O}(1)$ is uniformly bounded with respect to (x, y, K, n) . By using the definition (53) of the pseudo-distance δ_n , we get the relation

$$\begin{aligned} \dim(E_{(Kn-K, Kn]}) \delta_n(x, y)^2 \\ = 2K^d \int_{n-1}^n \nu^{d-1} [\widehat{\sigma}_{d-1}(0) - \widehat{\sigma}_{d-1}(K\nu \delta_g(x, y))] \frac{d\nu}{(2\pi)^{d/2}} + (Kn)^{d-1} \mathcal{O}(1). \end{aligned}$$

Remembering that the function $|\widehat{\sigma}_{d-1}|$ is bounded by $\widehat{\sigma}_{d-1}(0)$ (see (56)), we can restrict our analysis on $[n - 1/2, n]$ so that $\delta_n(x, y)^2$ is bounded from below by

$$(59) \quad \frac{2K^d}{\dim(E_{(Kn-K, Kn]})} \int_{n-1/2}^n \nu^{d-1} [\widehat{\sigma}_{d-1}(0) - \widehat{\sigma}_{d-1}(K\nu \delta_g(x, y))] \frac{d\nu}{(2\pi)^{d/2}} + \frac{(Kn)^{d-1} \mathcal{O}(1)}{\dim(E_{(Kn-K, Kn]})}.$$

By looking the last integral and our assumptions, one notices

$$\forall \nu \in [n - 1/2, n], \quad K\nu \delta_g(x, y) > (n - 1/2) \beta/n > \beta/2.$$

The inequality (49) (proved in the case $\mathcal{M} = S^d$) may be used to ensure the existence of $\rho \in (0, 1)$ (merely depending on the dimension d and on β) such that the following holds true

$$\forall \nu \in [n - 1/2, n], \quad \widehat{\sigma}_{d-1}(K\nu \delta_g(x, y)) \in \rho \widehat{\sigma}_{d-1}(0).$$

As at the end of the proof of Proposition 12, we obtain an inequality similar to (57) which in turn gives suitable constants K_0 and c .

PROPOSITION 14. — *There are three numbers $\beta_0 \in (0, 1)$, $K_0 > 1$ and $c > 1$ (depending on the Riemannian manifold \mathcal{M}) satisfying the following property: for any $K > K_0$, for any $n \in \mathbb{N}^*$ and any pair $(x, y) \in \mathcal{M}^2$, the following implication holds true*

$$\delta_g(x, y) < \frac{\beta_0}{Kn} \implies \frac{Kn \delta_g(x, y)}{c} \in \delta_n(x, y) \in cKn \delta_g(x, y).$$

Proof. — The main issue is that an inequality like (59) seems to give an unavoidable $\mathcal{O}(1)$ remainder. Forgetting technicalities, one may consider that the main idea relies on the formula (60) that will need derivatives of order 2 of the spectral function. In the next steps, we will make several restrictions so that any small enough number β_0 will be convenient.

Step 1. — We first recall that if β_0 is small enough, then any two points x and y of \mathcal{M} , satisfying $\delta_g(x, y) < \beta_0$, belong to a same geodesically convex Riemannian ball. To see that point, we invoke the Whitehead theorem (see [CE08, Th.5.14]): there is a continuous function $c : \mathcal{M} \rightarrow (0, +\infty)$, called the convexity radius, such that each Riemannian ball $B_{\delta_g}(z, r)$, for $z \in \mathcal{M}$, is geodesically convex for $r < c(z)$.

We finish by using the compactness of \mathcal{M} and choosing β_0 less than the positive number $\min_{z \in \mathcal{M}} c(z)$.

Step 2. — We now claim that if β_0 is small enough, then for any $z \in \mathcal{M}$ the open Riemannian ball $B_{\delta_g}(z, \beta_0)$ is relatively compact in the domain of a suitable local chart around z . We again give a compactness argument. Let $r(z) > 0$ be such that there is a coordinate patch around z (for some open subset U_z of \mathbb{R}^d)

$$\begin{array}{ccc} \mathbb{R}^d & & \mathcal{M} \\ \cup & & \cup \\ \tau_z : U_z & \longrightarrow & B_{\delta_g}(z, r(z)). \end{array}$$

By compactness, one may cover the compact manifold \mathcal{M} with a finite atlas as follows

$$\mathcal{M} = B_{\delta_g}(z_1, r(z_1)) \cup \cdots \cup B_{\delta_g}(z_\ell, r(z_\ell)).$$

In other words, any $z \in \mathcal{M}$ satisfies $\max_{1 \leq k \leq \ell} (r(z_k) - \delta_g(z, z_k)) > 0$. By continuity and compactness, one may enforce the previous inequality as follows for some positive constant R :

$$\max_{1 \leq k \leq \ell} (r(z_k) - \delta_g(z, z_k)) > 2R.$$

Hence, by choosing $\beta_0 \in R$, for any $z \in \mathcal{M}$, there is $k \in \{1, \dots, \ell\}$ such that we have the inclusion

$$\overline{B_{\delta_g}(z, \beta_0)} \subset \overline{B_{\delta_g}(z_k, r(z_k) - R)}.$$

In other words, $B_{\delta_g}(z, \beta_0)$ is relatively compact in the domain $B_{\delta_g}(z_k, r(z_k))$.

Step 3. — Now fix x and y satisfying $\delta_g(x, y) < \beta_0/Kn$ as in the statement (other restrictions on β_0 will be given in Step 4 and Step 5). Since we are looking for $K > 1$, we also have $\delta_g(x, y) < \beta_0$. Thanks to Step 2, we can consider a coordinate patch $\tau_k : U_{z_k} \subset \mathbb{R}^d \rightarrow B_{\delta_g}(z_k, r(z_k)) \subset \mathcal{M}$ so that the ball $B_{\delta_g}(x, \beta_0)$ is included in the compact set $\overline{B_{\delta_g}(z_k, r(z_k) - R)}$. Thanks to Step 1, there is a geodesic

$$\gamma : [0, 1] \longrightarrow \overline{B_{\delta_g}(z_k, r(z_k) - R)}$$

starting from x and stopping at y . By using the symmetry of $(x, y) \mapsto e_{(Kn-K, Kn]}(x, y)$, we obtain the following integral formulas

$$\begin{aligned} & e_{(Kn-K, Kn]}(x, x) + e_{(Kn-K, Kn]}(y, y) - 2e_{(Kn-K, Kn]}(x, y) \\ &= e_{(Kn-K, Kn]}(y, y) - e_{(Kn-K, Kn]}(y, x) \\ & \quad - [e_{(Kn-K, Kn]}(x, y) - e_{(Kn-K, Kn]}(x, x)] \\ (60) \quad &= \int_0^1 \frac{\partial}{\partial t_1} \{e_{(Kn-K, Kn]}(\gamma(t_1), y) - e_{(Kn-K, Kn]}(\gamma(t_1), x)\} dt_1 \\ &= \int_0^1 \int_0^1 \frac{\partial^2}{\partial t_1 \partial t_2} \{e_{(Kn-K, Kn]}(\gamma(t_1), \gamma(t_2))\} dt_1 dt_2. \end{aligned}$$

Step 4. — Assuming that β_0 is smaller than the constant α_0 of Theorem 11, it is natural to compare the previous integral formula by replacing $e_{(K_{n-K}, K_n]}$ with its principal term appearing in (58). More precisely, let us introduce

$$(61) \quad e_{(K_{n-K}, K_n]}^\varphi(x, y) = K^d \int_{n-1}^n \nu^{d-1} \mathfrak{F}_{d-1}(K\nu\delta_g(x, y)) \frac{d\nu}{(2\pi)^{d/2}}.$$

We now claim that the following inequality holds true for any $(t_1, t_2) \in [0, 1]^2$:

$$(62) \quad \left| \frac{\partial^2}{\partial t_1 \partial t_2} \left\{ e_{(K_{n-K}, K_n]}(\gamma(t_1), \gamma(t_2)) - e_{(K_{n-K}, K_n]}^\varphi(\gamma(t_1), \gamma(t_2)) \right\} \right| \leq C(Kn)^{d+1} \delta_g(x, y)^2,$$

where C merely depend on the Riemannian structure of \mathcal{M} . To obtain that bound, we work with the coordinate patch $\tau_k : U_{z_k} \subset \mathbb{R}^d \rightarrow B_{\delta_g}(z_k, r(z_k)) \subset \mathcal{M}$ around x and y as above:

$$e_{(K_{n-K}, K_n]}(\gamma(t_1), \gamma(t_2)) = \overbrace{e_{(K_{n-K}, K_n]} \circ (\tau_k \oplus \tau_k)}^{U_{z_k} \times U_{z_k} \rightarrow \mathbb{R}} \left(\tau_k^{-1} \circ \gamma(t_1), \tau_k^{-1} \circ \gamma(t_2) \right).$$

By denoting (p, q) a generic point of $U_{z_k} \times U_{z_k}$, the previous formula allows us to write the double derivative

$$\frac{\partial^2}{\partial t_1 \partial t_2} \left\{ e_{(K_{n-K}, K_n]}(\gamma(t_1), \gamma(t_2)) \right\}$$

as follows (in which we denote by $(\tau_k^{-1})_i$ the i -th coordinate of τ_k^{-1}):

$$\sum_{i=1}^d \sum_{j=1}^d \frac{d}{dt_1} \left((\tau_k^{-1})_i \circ \gamma(t_1) \right) \times \frac{d}{dt_2} \left((\tau_k^{-1})_j \circ \gamma(t_2) \right) \times \Lambda_{ij}$$

with $\Lambda_{ij} = \frac{\partial^2}{\partial p_i \partial q_j} \left\{ e_{(K_{n-K}, K_n]} \circ (\tau_k \oplus \tau_k) \right\} \Big|_{(\tau_k^{-1} \circ \gamma(t_1), \tau_k^{-1} \circ \gamma(t_2))}$.

A similar computation holds true for $e_{(K_{n-K}, K_n]}^\varphi$. Here is the point where we really need the asymptotic given by Theorem 11 of the spectral function with derivatives and holding uniformly on any compact subset of the domain $B_{\delta_g}(z_k, r(z_k))$ of the local chart τ_k^{-1} given by the coordinate patch τ_k . More precisely we choose the compact subset $\overline{B_{\delta_g}(z_k, r(z_k)) - R}$ that turns out to contain the geodesic γ (see Step 3). Theorem 11 then shows the bound (uniformly in $(t_1, t_2) \in [0, 1]^2$):

$$\frac{\partial^2}{\partial p_i \partial q_j} \left\{ (e_{(K_{n-K}, K_n]} - e_{(K_{n-K}, K_n]}^\varphi) \circ (\tau_k \oplus \tau_k) \right\} \Big|_{(\tau_k^{-1} \circ \gamma(t_1), \tau_k^{-1} \circ \gamma(t_2))} = \mathcal{O}((Kn)^{d+1}).$$

And hence we get

$$\left| \frac{\partial^2}{\partial t_1 \partial t_2} \left\{ e_{(K_{n-K}, K_n]}(\gamma(t_1), \gamma(t_2)) - e_{(K_{n-K}, K_n]}^\varphi(\gamma(t_1), \gamma(t_2)) \right\} \right| \leq C(Kn)^{d+1} \sum_{i=1}^d \sum_{j=1}^d \left| \frac{d}{dt_1} \left((\tau_k^{-1})_i \circ \gamma(t_1) \right) \right| \times \left| \frac{d}{dt_2} \left((\tau_k^{-1})_j \circ \gamma(t_2) \right) \right|.$$

where C merely depends on the Riemannian structure of \mathcal{M} (we recall that Step 2 allows us to work in a finite atlas and with a finite collection of compact subsets $\overline{B_{\delta_g}(z_k, r(z_k) - R)}$). Remembering now that $\gamma : [0, 1] \rightarrow \mathcal{M}$ is a geodesic between x and y , the speed of γ is constant and must equal $\delta_g(x, y)$ (since we have parametrized the curve with $[0, 1]$). In other words, we have $\|\gamma'(t_1)\|_{T_{\gamma(t_1)}\mathcal{M}} = \delta_g(x, y)$ in the tangent space $T_{\gamma(t_1)}\mathcal{M}$ endowed with its Riemannian inner product. Still using that we are working with a finite atlas, we clearly obtain, for a suitable uniform constant $C > 0$, the following bound

$$\forall t_1 \in [0, 1], \quad \left| \frac{d}{dt_1} ((\tau_k^{-1})_i \circ \gamma(t_1)) \right| \leq C \|\gamma'(t_1)\|_{T_{\gamma(t_1)}\mathcal{M}} = C\delta_g(x, y).$$

A similar reasoning may be done for t_2 and (62) is proved.

Step 5. — We now claim that if β_0 is small enough, then the contribution of the principal term $e_{(Kn-K, Kn]}^{\wp}$ defined in (61) is given by the following equivalence (uniformly in $(t_1, t_2) \in [0, 1]^2$):

$$(63) \quad \frac{\partial^2}{\partial t_1 \partial t_2} e_{(Kn-K, Kn]}^{\wp}(\gamma(t_1), \gamma(t_2)) \simeq K(Kn)^{d+1} \delta_g(x, y)^2,$$

where the constants of equivalence merely depend on the dimension $d = \dim(\mathcal{M})$. To prove such estimates, we shall use the equality $\delta_g(\gamma(t_1), \gamma(t_2)) = \delta_g(x, y)|t_1 - t_2|$ coming from the fact that geodesic $\gamma : [0, 1] \rightarrow \mathcal{M}$ has a speed equaling $\delta_g(x, y)$. Since the function $\mathfrak{F}_{d-1} : (0, +\infty) \rightarrow \mathbb{R}$ admits an even smooth extension to \mathbb{R} (see (48) and (56)), one may write

$$\begin{aligned} \mathfrak{F}_{d-1}(K\nu\delta_g(\gamma(t_1), \gamma(t_2))) &= \mathfrak{F}_{d-1}(K\nu\delta_g(x, y)|t_1 - t_2|) \\ &= \mathfrak{F}_{d-1}(K\nu\delta_g(x, y)(t_1 - t_2)) \end{aligned}$$

and hence

$$\begin{aligned} \frac{\partial^2}{\partial t_1 \partial t_2} e_{(Kn-K, Kn]}^{\wp}(\gamma(t_1), \gamma(t_2)) &= K^d \int_{n-1}^n \nu^{d-1} \frac{\partial^2}{\partial t_1 \partial t_2} \mathfrak{F}_{d-1}(K\nu\delta_g(x, y)(t_1 - t_2)) \frac{d\nu}{(2\pi)^{d/2}} \\ &= -K^{d+2} \delta_g(x, y)^2 \int_{n-1}^n \nu^{d+1} \mathfrak{F}_{d-1}''(K\nu\delta_g(x, y)(t_1 - t_2)) \frac{d\nu}{(2\pi)^{d/2}}. \end{aligned}$$

Remember now that x and y satisfy the inequality $\delta_g(x, y) < \beta_0/Kn$. In the last integral, we see that the term $K\nu\delta_g(x, y)(t_1 - t_2)$ belongs to the interval $(-\beta_0, \beta_0)$. Differentiating and taking the real part of the Poisson formula (48) give

$$-\mathfrak{F}_{d-1}''(t) = \frac{1}{2^{d/2-1} \Gamma((d-1)/2) \sqrt{\pi}} \int_{-1}^1 \cos(ts) s^2 (1-s^2)^{(d-3)/2} ds.$$

Here is our last restriction for β_0 : due to the continuity of \mathfrak{F}_{d-1}'' at 0, if β_0 is small enough then one obviously has

$$\exists c > 0, \exists C > 0, \forall t \in (-\beta_0, \beta_0), \quad c \leq -\mathfrak{F}_{d-1}''(t) \leq C.$$

These inequalities lead to (63).

Step 6. — By plugging (62) and (63) in (60), we get the following inequality (for suitable positive constants C_1 and C_2):

$$e_{(Kn-K,Kn]}(x, x) + e_{(Kn-K,Kn]}(y, y) - 2e_{(Kn-K,Kn]}(x, y) > (C_1K^{d+2} - C_2K^{d+1})n^{d+1}\delta_g(x, y)^2.$$

Coming back to (53) and (51), there is also another positive constant $C_3 > 0$ satisfying

$$\delta_n(x, y)^2 > \left[\frac{C_1K^{d+2} - C_2K^{d+1}}{C_3K^d} \right] n^2\delta_g(x, y)^2.$$

A similar argument would show

$$\delta_n(x, y)^2 \leq \left[\frac{C_1K^{d+2} + C_2K^{d+1}}{C_3K^d} \right] n^2\delta_g(x, y)^2.$$

We conclude by making K tend to $+\infty$.

Theorem 10 follows by an application of Proposition 13 for the constant $\beta = \beta_0 \in (0, 1)$ of Proposition 14. Upon increasing the constants K_0 and c , we assume that those numbers have the same meaning in Proposition 13 and Proposition 14. In the zone $\delta_g > \beta_0/Kn$, we have $1/c \leq \delta_n \leq c$ and hence

$$\frac{\beta_0}{c} \min(1, Kn\delta_g) \leq \delta_n \leq \frac{c}{\beta_0} \min(1, Kn\delta_g).$$

Similarly, in the zone $\delta_g < \beta_0/Kn$, the inequalities $(1/c)Kn\delta_g \leq \delta_n \leq cKn\delta_g$ hold true and so do the following ones

$$\frac{1}{c} \min(1, Kn\delta_g) \leq \delta_n \leq c \min(1, Kn\delta_g).$$

Hence we get (55) in which one may choose $C = c/\beta_0$.

7. PRELIMINARIES FOR PROOFS OF THEOREMS 2 AND 3

For our purpose, a Gaussian process $(f^{G,\omega}(x))_{x \in \mathcal{M}}$ on (\mathcal{M}, δ_g) is defined as a family of real random variables satisfying the following two properties

(i) for any finite subset $\{x_1, \dots, x_k\} \subset \mathcal{M}$, the random vector

$$\omega \in \Omega \longmapsto (f^{G,\omega}(x_1), \dots, f^{G,\omega}(x_k)) \in \mathbb{R}^k$$

is a centered Gaussian vector. In other words, any linear combination

$$\omega \longmapsto \sum_{i=1}^k \alpha_i f^{G,\omega}(x_i),$$

with $\alpha_i \in \mathbb{R}$, is centered and Gaussian.

(ii) the function $(x, y) \in \mathcal{M}^2 \mapsto \mathbf{E}_\omega[f^{G,\omega}(x)f^{G,\omega}(y)]$, called the covariance structure, is continuous. An equivalent condition is the continuity of the function $x \mapsto f^{G,\omega}(x)$ from \mathcal{M} to $L^2(\Omega)$.

Here is the simple example of a Gaussian process we are interested in: for any continuous function $c : \mathcal{M} \rightarrow \ell^2(\mathbb{N})$ the process $(f^{G,\omega}(x))_{x \in \mathcal{M}}$ given by $f^{G,\omega}(x) = \sum_{\ell \in \mathbb{N}} g_\ell(\omega) c_\ell(x)$ is Gaussian. As recalled in the introduction, we shall need the following pseudo-distance δ given by

$$\forall (x, y) \in \mathcal{M}^2, \quad \delta(x, y) := \sqrt{\mathbf{E}_\omega[|f^{G,\omega}(x) - f^{G,\omega}(y)|^2]}.$$

We note that if the Gaussian random series $\sum f_n^{G,\omega}$ defined in (21) almost surely converges in $\mathcal{C}^0(\mathcal{M}_s)$ (for a submanifold \mathcal{M}_s of \mathcal{M}) then the random series $\sum f_n^{G,\omega}(x)$ almost surely converges for any $x \in \mathcal{M}_s$. For the sake of clarity, we state the following result.

PROPOSITION 15. — *There is a constant $K_0 > 1$ merely depending on the Riemannian manifold \mathcal{M} such that for any $K > K_0$ and for any sequence $(f_n)_{n>1}$ satisfying $f_n \in E_{(K_n-K, K_n]}$ for $n > 1$ (see (13)), then the following assertions are equivalent:*

- (i) *the series $\sum \|f_n\|_{L^2(\mathcal{M})}^2$ is convergent,*
- (ii) *for any $x \in \mathcal{M}$, the random series $\sum f_n^{G,\omega}(x)$ converges in $L^2(\Omega)$ to a Gaussian random variable,*
- (iii) *for any $x \in \mathcal{M}$, the random series $\sum f_n^{G,\omega}(x)$ almost surely converges in \mathbb{R} ,*
- (iv) *there is $x \in \mathcal{M}$ such that the random series $\sum f_n^{G,\omega}(x)$ almost surely converges in \mathbb{R} .*

Moreover, the previous statements imply the following one:

(v) *the family of random variables $f^{G,\omega}(x) = \sum_{n>1} f_n^{G,\omega}(x)$ is a Gaussian process on \mathcal{M} and its Dudley pseudo-distance $\delta : \mathcal{M}^2 \rightarrow [0, +\infty)$ is given, for any $(x, y) \in \mathcal{M}^2$, by*

$$(64) \quad \delta(x, y)^2 = \mathbf{E}_\omega[|f^{G,\omega}(x) - f^{G,\omega}(y)|^2] = \sum_{n>1} \|f_n\|_{L^2(\mathcal{M})}^2 \delta_n(x, y)^2,$$

where the partial pseudo-distance δ_n is defined in (52).

Finally, in the specific case $\mathcal{M} = S^d$, a similar statement holds true if each f_n belongs to the eigenspace $\ker(\Delta + n(n+d-1))$ provided that we replace $f_n^{G,\omega}$ with (20) and δ_n with (35).

Proof

(i) \Leftrightarrow (ii). — For any $x \in \mathcal{M}$, the random variable $f_n^{G,\omega}(x)$ is centered and Gaussian with variance equaling

$$\mathbf{E}_\omega[|f_n^{G,\omega}(x)|^2] = \frac{\|f_n\|_{L^2(\mathcal{M})}^2}{\dim(E_{(K_n-K, K_n]})} \sum_{\lambda_k \in (K_n-K, K_n]} \phi_k(x)^2.$$

By orthogonality of the random variables $(f_n^{G,\omega}(x))_{n>1}$ in $L^2(\Omega)$, it is clear and well-known that the Gaussian series $\sum f_n^{G,\omega}(x)$ converges in $L^2(\Omega)$ if and only if the series $\sum \mathbf{E}_\omega[|f_n^{G,\omega}(x)|^2]$ converges and we have

$$\mathbf{E}_\omega \left[\left| \sum_{n>1} f_n^{G,\omega}(x) \right|^2 \right] = \sum_{n>1} \mathbf{E}_\omega[|f_n^{G,\omega}(x)|^2].$$

Thanks to (50) and (51), we get the equivalence (i) \Leftrightarrow (ii) and there are two constants $C > 1$ and $K_0 > 1$ that merely depend on the Riemannian structure of \mathcal{M} such that for any $K > K_0$ we have

$$(65) \quad \frac{1}{C} \sum_{n>1} \|f_n\|_{L^2(\mathcal{M})}^2 \preceq \mathbf{E}_\omega \left[\left| \sum_{n>1} f_n^{G,\omega}(x) \right|^2 \right] \preceq C \sum_{n>1} \|f_n\|_{L^2(\mathcal{M})}^2.$$

(iii) \Leftrightarrow (iv). — Obvious.

(i) \Leftrightarrow (iii) and (iv) \Leftrightarrow (i). — See for instance [LQ18b, Cor. III.6, p. 26] and the previous computations.

Proof of (v). — The finite linear combination $\sum_{i=1}^p \alpha_i (\sum_{n>1} f_n^{G,\omega}(x_i))$, with $\alpha_i \in \mathbb{R}$, is Gaussian because it is the limit in $L^2(\Omega)$ of the Gaussian random variables

$$\sum_{n=1}^N \left(\sum_{i=1}^p \alpha_i f_n^{G,\omega}(x_i) \right)$$

as $N \rightarrow +\infty$ (with the same argument as in (i) \Leftrightarrow (iii)).

We now have to check the continuity of the covariance structure. By using the notation (52) and orthogonality arguments, the definition (21) of $f_n^{G,\omega}$ directly proves (64). We recall that the expected continuity is equivalent to the continuity of the function $x \in \mathcal{M} \mapsto f^{G,\omega}(x) \in L^2(\Omega)$. But the partial pseudo-distance δ_n are continuous on \mathcal{M}^2 and uniformly bounded on \mathcal{M}^2 (see (54)). Hence, the right-hand side of (64) absolutely converges in $\mathcal{C}^0(\mathcal{M}^2)$ and we obtain the expected continuity.

Towards the case $\mathcal{M} = S^d$. — For the sphere S^d , we work with (20). In the same spirit as (38), we have

$$\sum_{\lambda_k^2 = n(n+d-1)} \phi_k(x)^2 = \frac{\dim(E_n)}{\text{vol}_{S^d}(S^d)}.$$

From such an identity, we see that the equivalence (65) becomes

$$\mathbf{E}_\omega \left[\left| \sum_{n>1} f_n^{G,\omega}(x) \right|^2 \right] = \frac{1}{\text{vol}_{S^d}(S^d)} \sum_{n>1} \|f_n\|_{L^2(S^d)}^2.$$

We also deduce that the partial pseudo-distances δ_n , associated to the sequence of the eigenspaces of Δ and defined in (35), are uniformly bounded on $S^d \times S^d$ (an information used in the proof of (v)):

$$(66) \quad \delta_n \preceq \frac{2}{\sqrt{\text{vol}_{S^d}(S^d)}}.$$

8. PROOF OF THEOREM 2 FOR SPHERES S^d WITH $d > 2$

Thanks to the convergence of $\sum_{n>1} \|f_n\|_{L^2(S^d)}^2$ and Point v) of Proposition 15, we know that the process $(\sum_{n>1} f_n^{G,\omega}(x))_{x \in S^d}$ is Gaussian and so is its restriction to any compact submanifold \mathcal{M}_s . We begin by the following proposition.

PROPOSITION 16. — *Let us consider a Gaussian process $(F^\omega(x))_{x \in S^d}$ whose covariance structure reads*

$$(67) \quad \forall (x, y) \in S^d \times S^d, \quad \mathbf{E}_\omega[F^\omega(x)F^\omega(y)] = C(\langle x, y \rangle)$$

for some continuous function $C : [-1, 1] \rightarrow \mathbb{R}$. Then the following statements are equivalent:

- (i) the condition $\int_0^1 ((C(1) - C(\cos(t)))^{1/2} / t \sqrt{\ln(1/t)}) dt < +\infty$ is fulfilled,
- (ii) the Gaussian process $(F^\omega(x))_{x \in S^d}$ admits a version $(\tilde{F}^\omega(x))_{x \in S^d}$ which is sample-continuous in the following sense:
 - with probability one, the function $x \mapsto \tilde{F}^\omega(x)$ is continuous from S^d to \mathbb{R} ,
 - for any $x \in \mathcal{M}$, the equality $\mathbf{P}[F^\omega(x) = \tilde{F}^\omega(x)] = 1$ holds true.
- (iii) on some fixed geodesic arc Γ of S^d , the Gaussian process $(F^\omega(x))_{x \in \Gamma}$ process on Γ admits a version which is sample-bounded in the following sense:
 - with probability 1, the function $x \mapsto \tilde{F}^\omega(x)$ is bounded from Γ to \mathbb{R} ,
 - for any $x \in \mathcal{M}$, the equality $\mathbf{P}[F^\omega(x) = \tilde{F}^\omega(x)] = 1$ holds true.

Proof

Preliminary remarks. — Note that due to the definition (67) of the function C and to the Cauchy-Schwarz inequality, we easily check the inequality $|C(u)| \leq C(1)$ for any $u \in [-1, 1]$ (so the square root in the integral in Point i) is well defined). Then one may compute the Dudley pseudo-distance δ of the Gaussian process $(F^\omega(x))_{x \in S^d}$:

$$\delta(x, y) := \sqrt{\mathbf{E}_\omega[|F^\omega(x) - F^\omega(y)|^2]} = \sqrt{2C(1) - 2C(\langle x, y \rangle)}.$$

We now note that for any two points x and y on the sphere S^d we have $\langle x, y \rangle = \cos(\delta_g(x, y))$ where $\delta_g(x, y)$ stands for the Riemannian distance on S^d . In other words, we have

$$(68) \quad \delta(x, y) = \sqrt{2C(1) - 2C(\cos(\delta_g(x, y)))}.$$

Now from the classical result [Sch42, p.101, Th.1] by Schoenberg, there exists a sequence of non-negative coefficients $(u_n)_{n > 0}$ satisfying

$$\forall \theta \in \mathbb{R}, \quad C(\cos(\theta)) = \sum_{n=0}^{+\infty} u_n P_n^{(d/2-1, d/2-1)}(\cos(\theta))$$

and such that series converges at $\theta = 0$ (in other words $\sum_{n=0}^{+\infty} n^{d/2-1} u_n < +\infty$).

(i) \Rightarrow (ii). — We may cover S^d by a finite set of closed Riemannian balls:

$$S^d = \bigcup_{1 \leq k \leq N} \overline{B_k} \quad \text{with } B_k := B_{\delta_g}(a_k, \pi/4) \text{ and } a_k \in S^d.$$

In order to show (ii), it is clear that it is sufficient to prove that each Gaussian process $(F^\omega(x))_{x \in \overline{B_k}}$, for $1 \leq k \leq N$, admits a sample-continuous version. Thanks to the Dudley theorem, it is sufficient to show the entropy conditions of the Dudley pseudo-distances of the N Gaussian processes $(F^\omega(x))_{x \in \overline{B_k}}$. Since each closed ball $\overline{B_k}$

has a diameter equaling $\pi/2 < \pi$, we may use Proposition 9 (with $\vartheta = \pi/4$) to get the equivalence (up to multiplicative constants independent of x and y):

$$\delta(x, y)^2 = 2 \sum_{n=1}^{+\infty} u_n \left[P_n^{(d/2-1, d/2-1)}(1) - P_n^{(d/2-1, d/2-1)}(\cos(\delta_g(x, y))) \right]$$

$$(69) \quad \delta(x, y)^2 \simeq \sum_{n=1}^{+\infty} u_n n^{d/2-1} \min(1, n\delta_g(x, y))^2.$$

The function $\Upsilon : t \in [0, +\infty) \mapsto \sqrt{\sum_{n>1} u_n n^{d/2-1} \min(1, nt)^2}$ fulfills the assumption of Proposition 6. Note that the probability Riemannian volume μ of S^d satisfies (28) (with suitable possibly different constants) on the ball $\overline{B_k}$ since $\overline{B_k}$ is a submanifold with smooth boundary. Then the conclusion of Proposition 6 combined with (68) and (69) shows that the entropy condition of each $(F^\omega(x))_{x \in \overline{B_k}}$ is equivalent to Point (i).

(ii) \Rightarrow (iii). — Obvious.

(iii) \Rightarrow (i). — The main point is that (67) implies that the restriction of the Gaussian process $(F^\omega(x))_{x \in S^d}$ is stationary on any geodesic closed curve (identified as the torus \mathbb{T}). By applying a finite set of translations to the Gaussian process $(F^\omega(x))_{x \in \Gamma}$, one obtains the almost sure boundedness on the whole geodesic curve containing Γ . So there is no loss of generality to assume that Γ is the whole geodesic curve of the points $x_\theta = (\cos(\theta), \sin(\theta), 0, \dots, 0)$ for $\theta \in [0, 2\pi]$. But the Dudley-Fernique theorem ensures that the almost sure boundedness is equivalent to the entropy condition on Γ . In order to use Proposition 6 and the equivalence (69) of the Dudley pseudo-distance, we need to restrict to a subcurve of Γ on which the geodesic distance is strictly less than π , for instance

$$(70) \quad \Gamma^+ := \{(\cos(\theta), \sin(\theta), 0, \dots, 0), \quad \theta \in [0, \pi/2]\}.$$

It is easy to see that the entropy condition on Γ^+ is still satisfied.⁽⁷⁾ Let us now denote by δ_{Γ^+} the Dudley pseudo-distance on Γ^+ of the Gaussian process $(F^\omega(x))_{x \in \Gamma^+}$. Due to the equivalence (69), we still have $\delta_{\Gamma^+} \simeq \Upsilon(\delta_g)$ (where δ_g equals both the geodesic distance on Γ^+ and of course the geodesic distance on S^d due to the definition of Γ^+). Moreover, the length measure of Γ^+ obviously satisfies inequalities like (28) (with $\sigma = 1$). As above, the conclusion of Proposition 6 shows Point (i).

In the sequel of this part, we shall consider the case of the Gaussian process $f^{G,\omega} = \sum_{n>1} f_n^{G,\omega}$ defined in (20) with $\sum_{n>1} \|f_n\|_{L^2(S^d)}^2 < +\infty$. The proof of Proposition 15

⁽⁷⁾We indeed have the inequality $N_\delta(\Gamma^+, 2\varepsilon) \leq N_\delta(\Gamma, \varepsilon)$. For the proof, we consider a finite cover $\Gamma = \bigcup B_\delta(x_i, \varepsilon)$ with $x_i \in \Gamma$. And so we have $\Gamma^+ = \bigcup (B_\delta(x_i, \varepsilon) \cap \Gamma^+)$. If $B_\delta(x_i, \varepsilon)$ intersects Γ^+ then any element $x'_i \in B_\delta(x_i, \varepsilon) \cap \Gamma^+$ satisfies $B_\delta(x_i, \varepsilon) \subset B_\delta(x'_i, 2\varepsilon)$. Hence we obtain a cover $\Gamma^+ \subset \bigcup B_\delta(x'_i, 2\varepsilon)$.

easily gives the formula

$$\forall (x, y) \in S^d \times S^d, \quad \mathbf{E}[f^{G,\omega}(x)f^{G,\omega}(y)] = \sum_{n>1} \frac{\|f_n\|_{L^2(S^d)}^2}{\dim(E_n)} \sum_{\lambda_k^2 = n(n+d-1)} \phi_k(x)\phi_k(y).$$

Hence, (38) shows that the previous term equals $C(\langle x, y \rangle)$ with

$$C(\theta) = \sum_{n>1} \frac{\|f_n\|_{L^2(S^d)}^2}{\dim(E_n)} c_{d,n} P_n^{(d/2-1, d/2-1)}(\cos(\theta))$$

(where $c_{d,n}$ is given by (39) and satisfies $c_{d,n} \simeq_{n \rightarrow +\infty} n^{d/2}$ and $\dim(E_n) \simeq_{n \rightarrow +\infty} n^{d-1}$) and the function $\Upsilon : [0, +\infty[\rightarrow [0, +\infty[$ of the previous proof can be chosen as

$$\Upsilon(t) = \sqrt{\sum_{n>1} \|f_n\|_{L^2(S^d)}^2 \min(1, nt)^2}.$$

Moreover, the Dudley pseudo-distance of $(f^{G,\omega}(x))_{x \in S^d}$ satisfies

$$(71) \quad \delta(x, y) \simeq \Upsilon(\delta_g(x, y))$$

uniformly in x and y provided that $\delta_g(x, y) \in \pi/2$ (that is a consequence of Theorem 8 and (64)). We now prove Theorem 2 and so we are interested in studying the almost sure continuity of $x \in \mathcal{M}_s \mapsto \sum_{n>1} f_n^{G,\omega}(x) \in \mathbb{R}$.

PROPOSITION 17. — *The equivalence (i) \Leftrightarrow (ii) of Theorem 2 is true.*

Proof

Preliminary remarks. — Let us recall why the existence of a sample-continuous version of the Gaussian process $(\sum_{n>1} f_n^{G,\omega}(x))_{x \in \mathcal{M}_s}$ on \mathcal{M}_s is equivalent to the almost sure convergence of the Gaussian random series $\sum f_n^{G,\omega}$ in $\mathcal{C}^0(\mathcal{M}_s)$, namely Point (i) of Theorem 2. That is the same argument than that the one used in the classical theory for compact groups. Actually, we apply the Itô-Nisio theorem (see [LQ18b, p. 238]) that merely needs to check the additional assumption ensuring that, for any $x \in \mathcal{M}_s$, each random numerical series $\sum f_n^{G,\omega}(x)$ is almost surely convergent. Such a fact is actually proved in Point iii) of Proposition 15.

Step 1

For (i) \Rightarrow (ii), we may clearly assume that \mathcal{M}_s is a curve with ends. We can also assume that the diameter of \mathcal{M}_s is less than $\pi/2$. Let $m_s : [0, \pi/2] \rightarrow \mathcal{M}_s$ be a bi-Lipschitz parametrization and let us consider the Gaussian process $(f^{G,\omega}(m_s(\theta)))_{\theta \in [0, \pi/2]}$. We clearly have the equivalence

$$\mathbf{E}[|f^{G,\omega}(m_s(\theta)) - f^{G,\omega}(m_s(\theta'))|^2] \simeq \Upsilon(\delta_g(m_s(\theta), m_s(\theta')))^2 \simeq \Upsilon(|\theta - \theta'|)^2.$$

Now let $\gamma : [0, \pi/2] \rightarrow \Gamma^+$ be the natural parametrization of Γ^+ defined in (70). We also have (see (71)):

$$(72) \quad \mathbf{E}[|f^{G,\omega}(\gamma(\theta)) - f^{G,\omega}(\gamma(\theta'))|^2] \simeq \Upsilon(|\theta - \theta'|)^2.$$

The classical Slepian comparison theorem ensures that the almost sure boundedness of the Gaussian processes $(f^{G,\omega}(m_s(\theta)))_{\theta \in [0, \pi/2]}$ is equivalent to that of

$(f^{G,\omega}(\gamma(\theta)))_{\theta \in [0, \pi/2]}$ (see [LT91, Cor. 3.14] or [LQ18b, p. 73]), and so the same is obviously true for $(f^{G,\omega}(x))_{x \in \mathcal{M}_s}$ and $(f^{G,\omega}(x))_{x \in \Gamma^+}$. Then Proposition 16 (or its proof) implies the following convergence

$$(73) \quad \int_0^1 \frac{\Upsilon(t)}{t\sqrt{\ln(1/t)}} dt < +\infty.$$

For the conclusion, we need to admit for a moment Step 2.

(i) \Rightarrow (ii). — That is a direct consequence of Step 2.

(ii) \Rightarrow (i). — By Step 2 we obtain (73), which is equivalent to Point i) of Proposition 16 (via (71)) which in turns proves the almost sure continuity of $x \mapsto f^{G,\omega}(x)$ (see the preliminary remarks above) on the whole sphere S^d thanks to Point (ii) of Proposition 16.

Step 2. — We shall prove that (73) is equivalent to the Salem-Zygmund condition (25). We begin by writing

$$\begin{aligned} \int_0^1 \sqrt{\sum_{n>1} \|f_n\|_{L^2(S^d)}^2 \min(1, n^2 t^2)} \frac{dt}{t\sqrt{-\ln(t)}} \\ = \sum_{p>1} \int_{1/(p+1)}^{1/p} \sqrt{\sum_{n>1} \|f_n\|_{L^2(S^d)}^2 \min(1, n^2 t^2)} \frac{dt}{t\sqrt{-\ln(t)}} \\ \simeq \sum_{p>1} \sqrt{\sum_{n>1} \|f_n\|_{L^2(S^d)}^2 \min(1, n^2/p^2)} \frac{1}{p\sqrt{\ln(p+1)}}. \end{aligned}$$

We now write

$$\sqrt{\sum_{n>1} \|f_n\|_{L^2(S^d)}^2 \min(1, n^2/p^2)} = \sqrt{\frac{1}{p^2} U_p + V_p} \simeq \frac{\sqrt{U_p}}{p} + \sqrt{V_p},$$

where U_p and V_p are defined as follows

$$(74) \quad U_p = \sum_{n=1}^{p-1} n^2 \|f_n\|_{L^2(S^d)}^2 \quad \text{and} \quad V_p = \sum_{n=p}^{+\infty} \|f_n\|_{L^2(S^d)}^2,$$

with the convention $U_1 = 0$. Hence we get

$$\int_0^1 \frac{\Upsilon(t)}{t\sqrt{-\ln(t)}} dt \simeq \sum_{p>1} \frac{\sqrt{U_p}}{p^2\sqrt{\ln(p+1)}} + \frac{\sqrt{V_p}}{p\sqrt{\ln(p+1)}}.$$

The V_p part is exactly the Salem-Zygmund type term in (25). So the proof will be finished provided that we show that the contribution of the U_p part is controlled by that of the V_p part:

$$(75) \quad \sum_{p>2} \frac{\sqrt{U_p}}{p^2\sqrt{\ln(p+1)}} \cdot \sum_{p>1} \frac{\sqrt{V_p}}{p\sqrt{\ln(p+1)}}.$$

Thanks to the Cauchy-Schwarz inequality and the definition of U_p in (74), we have

$$\begin{aligned} \sum_{p>2} \frac{\sqrt{U_p}}{p^2 \sqrt{\ln(p+1)}} &= \sum_{p>2} \frac{1}{\sqrt{p} \ln(p+1)} \times \frac{\sqrt{U_p} \sqrt{\ln(p+1)}}{p^{3/2}} \\ &\leq C \left(\sum_{p>2} \frac{U_p \ln(p+1)}{p^3} \right)^{1/2} \\ &\leq C \left(\sum_{n>1} \|f_n\|_{L^2(S^d)}^2 n^2 \sum_{p>n+1} \frac{\ln(p+1)}{p^3} \right)^{1/2} \\ &\leq C \left(\sum_{n=1}^{+\infty} \|f_n\|_{L^2(\mathcal{M})}^2 \ln(n+1) \right)^{1/2}. \end{aligned}$$

By using the equivalence $\sqrt{\ln(n+1)} \simeq \sum_{p=1}^n 1/p \sqrt{\ln(p+1)}$ and the definition of V_p in (74), we obtain

$$\begin{aligned} \sum_{p>1} \frac{\sqrt{U_p}}{p^2 \sqrt{\ln(p+1)}} &\leq C \left(\sum_{n=1}^{+\infty} \|f_n\|_{L^2(S^d)}^2 \left(\sum_{p=1}^n \frac{1}{p \sqrt{\ln(p+1)}} \right)^2 \right)^{1/2} \\ &\leq C \left(\sum_{n=1}^{+\infty} \|f_n\|_{L^2(S^d)}^2 \sum_{p=1}^n \sum_{q=1}^n \frac{1}{p \sqrt{\ln(p+1)} q \sqrt{\ln(q+1)}} \right)^{1/2} \\ &\leq C \left(\sum_{p=1}^{+\infty} \sum_{q=1}^{+\infty} \frac{V_{\max(p,q)}}{p \sqrt{\ln(p+1)} q \sqrt{\ln(q+1)}} \right)^{1/2}. \end{aligned}$$

By bounding $V_{\max(p,q)} \leq \sqrt{V_p V_q}$, we finally prove (75).

The following proposition finishes the proof of Theorem 2 for spheres.

PROPOSITION 18. — *The equivalence (i) \Leftrightarrow (iii) is true in Theorem 2.*

Proof. — The implication (iii) \Rightarrow (i) is nothing other than the Dudley theorem. Let us prove the converse implication (i) \Rightarrow (iii). Let us set $\sigma = \dim(\mathcal{M}_s)$. By compactness, we can decompose \mathcal{M}_s as a union of σ -dimensional compact submanifolds $\mathcal{M}_{s,1}, \dots, \mathcal{M}_{s,N}$ with smooth boundary and with diameter less than $\pi/2$:

$$(76) \quad \mathcal{M}_s = \mathcal{M}_{s,1} \cup \dots \cup \mathcal{M}_{s,N}, \quad \max_{\substack{x \in \mathcal{M}_{s,k} \\ y \in \mathcal{M}_{s,k}}} \delta_g(x, y) \leq \frac{\pi}{2}, \quad \forall k \in \{1, \dots, N\}.$$

Point (i) of Theorem 2 implies the almost sure continuity of each $x \in \mathcal{M}_{s,k} \mapsto f^{G,\omega}(x)$ for any $k \in \{1, \dots, N\}$. Proposition 17 (or its proof) shows that the following condition holds

$$\int_0^1 \frac{\Upsilon(t)}{t \sqrt{\ln(1/t)}} dt < +\infty.$$

Note that the probability Riemannian volume of each $\mathcal{M}_{s,k}$ satisfies the assumption (28). As a consequence of Proposition 6, we obtain the entropy condition for

each $\mathcal{M}_{s,k}$:

$$\int_0^{+\infty} \sqrt{\ln(N_\delta(\mathcal{M}_{s,k}, \varepsilon))} d\varepsilon < +\infty.$$

The definition of the covering number and (76) lead to the following inequality

$$(77) \quad N_\delta(\mathcal{M}_s, \varepsilon) \subset \sum_{1 \subset k \subset N} N_\delta(\mathcal{M}_{s,k}, \varepsilon),$$

which in turn implies

$$(78) \quad N_\delta(\mathcal{M}_s, \varepsilon) \subset N \times \prod_{1 \subset k \subset N} N_\delta(\mathcal{M}_{s,k}, \varepsilon).$$

We then easily deduce Point (iii) of Theorem 2.

9. PROOF OF THEOREM 3 FOR S^d WITH $d > 2$

The proof developed in the last part already shows that the following two numbers are equivalent (up to multiplicative constants):

- (i) the number $\int_0^1 (\Upsilon(t)/t\sqrt{\ln(1/t)}) dt$,
- (ii) the Salem-Zygmund expression $\sum_{p>1} (1/p\sqrt{\ln(p+1)}) (\sum_{n>p} \|f_n\|_{L^2(S^d)}^2)^{1/2}$.

In order to add the the entropy integral $\int_0^{+\infty} \sqrt{\ln(N_\delta(\mathcal{M}_s, \varepsilon))} d\varepsilon$ to that list, we merely need the following two arguments.

(a) The entropy integral on \mathcal{M}_s is equivalent to $\sup_{1 \subset k \subset N} \int_0^{+\infty} \sqrt{\ln(N_\delta(\mathcal{M}_{s,k}, \varepsilon))} d\varepsilon$. To see that equivalence, one firstly precises (77) as follows

$$\max_{1 \subset k \subset N} N_\delta(\mathcal{M}_{s,k}, 2\varepsilon) \subset N_\delta(\mathcal{M}_s, \varepsilon),$$

which can be proved as explained after (70). Then, we easily can bound from below the entropy integral $\int_0^{+\infty} \sqrt{\ln(N_\delta(\mathcal{M}_s, \varepsilon))} d\varepsilon$. To bound from above the entropy integral on \mathcal{M}_s by $\sup_{1 \subset k \subset N} \int_0^{+\infty} \sqrt{\ln(N_\delta(\mathcal{M}_{s,k}, \varepsilon))} d\varepsilon$, we may use (78) by remembering that N in (78) can be chosen merely depending on the submanifold \mathcal{M}_s and that the diameter of (\mathcal{M}_s, δ) is bounded by $C(d)\sqrt{\lim_{n>1} \|f_n\|_{L^2(S^d)}^2}$ (see (66)).

(b) Thanks to Corollary 7, each entropy integral $\int_0^{+\infty} \sqrt{\ln(N_\delta(\mathcal{M}_{s,k}, \varepsilon))} d\varepsilon$ of $\mathcal{M}_{s,k}$ is equivalent to the same number $\int_0^1 (\Upsilon(t)/t\sqrt{\ln(1/t)}) dt$.

The last thing to explain is why we can equivalently control the expectation

$$\mathbf{E}_\omega \left[\sup_{x \in \mathcal{M}_s} \left| \sum_{n>1} f_n^{G,\omega}(x) \right| \right].$$

The quantitative version of the Dudley theorem will allow to bound from above the last expectation by the entropy integral (see [LT91, Th. 11.17]): we fix $x_0 \in \mathcal{M}_s$ and

write (thanks to the symmetry of the Gaussian process):

$$\begin{aligned} \mathbf{E}_\omega \left[\sup_{x \in \mathcal{M}_s} \left| \sum_{n>1} f_n^{G,\omega}(x) \right| \right] &\leq \mathbf{E}_\omega \left[\left| \sum_{n>1} f_n^{G,\omega}(x_0) \right| \right] + \mathbf{E}_\omega \left[\sup_{x,y \in \mathcal{M}_s} \left| \sum_{n>1} f_n^{G,\omega}(x) - f_n^{G,\omega}(y) \right| \right] \\ &\leq \mathbf{E}_\omega \left[\left| \sum_{n>1} f_n^{G,\omega}(x_0) \right| \right] + 2\mathbf{E}_\omega \left[\sup_{x \in \mathcal{M}_s} \sum_{n>1} f_n^{G,\omega}(x) \right]. \end{aligned}$$

The last expectation is controlled by the entropy integral ([LT91, Th. 11.17]) whereas the first is controlled as follows (see the proof of Proposition 15):

$$\begin{aligned} \mathbf{E}_\omega \left[\left| \sum_{n>1} f_n^{G,\omega}(x_0) \right| \right] &\leq \mathbf{E}_\omega \left[\left| \sum_{n>1} f_n^{G,\omega}(x_0) \right| \right]^{1/2} \leq C \left(\sum_{n>1} \|f_n\|_{L^2(S^d)}^2 \right)^{1/2} \\ &\leq C \sqrt{\ln(2)} \sum_{p>1} \frac{1}{p\sqrt{\ln(p+1)}} \left(\sum_{n>p} \|f_n\|_{L^2(S^d)}^2 \right)^{1/2} \quad (\text{see } p=1) \end{aligned}$$

and then we note that we already proved that the Salem-Zygmund term is controlled by the entropy integral at the beginning of this part.

For bounding $\mathbf{E}_\omega [\sup_{x \in \mathcal{M}_s} |\sum_{n>1} f_n^{G,\omega}(x)|_{\mathcal{C}^0(\mathcal{M}_s)}]$ from below, we may use the quantitative version of the Fernique theorem (see [LT91, Th. 13.3]) and the Slepian comparison theorem (see the proofs of (iii) \Rightarrow (i) in Proposition 16 and Step 1 in Proposition 17).

10. PROOF OF THEOREM 2 AND 3 FOR COMPACT MANIFOLDS

Here $f^{G,\omega}$ is defined in (21) and so the Dudley pseudo-distance δ implicitly depends on K since $f_n^{G,\omega}$ is a random wave with values in $E_{(K_n-K, K_n)}$. We assume that K is large enough so that the conclusion of Theorem 10 holds true.

– For the proof of the analogue of Proposition 17, the mere thing to clarify is the proof of Step 1. Actually, we can repeat some arguments with

$$\Upsilon(t) = \sqrt{\sum_{n>1} \|f_n\|_{L^2(\mathcal{M})}^2 \min(1, Knt)^2} \simeq \sqrt{\sum_{n>1} \|f_n\|_{L^2(\mathcal{M})}^2 \min(1, nt)^2}.$$

Note the equivalence $\delta \simeq \Upsilon(\delta_g)$. For (i) \Rightarrow (ii), the idea⁽⁸⁾ is that the Slepian theorem shows that the following two Gaussian processes are simultaneously sample-continuous or simultaneously unbounded:

- $(f^{G,\omega}(x))_{x \in \mathcal{M}_s}$ on a curve \mathcal{M}_s of a compact manifold \mathcal{M} and $f^{G,\omega}(x)$ defined in (21),
- $(f^{G,\omega}(\gamma(\theta)))_{\theta \in [0, \pi/2]}$ with $\gamma : [0, \pi/2] \rightarrow \Gamma^+$ as in (72) but $(f^{G,\omega}(x))_{x \in S^d}$ defined on the sphere S^d (as in (20) with $\|f_n\|_{L^2(\mathcal{M})}$ instead of $\|f_n\|_{L^2(S^d)}$) with

⁽⁸⁾The author is grateful to a referee for that argument.

the following form:

$$f^{G,\omega}(x) := \sum_{n>1} \frac{\|f_n\|_{L^2(\mathcal{M})}}{\sqrt{\dim(E_n)}} \sum_{\lambda_k^2=n(n+d-1)} g_{n,k}(\omega)\phi_k(x), \quad x \in S^d.$$

Repeating the end of argument of Step 1 of Proposition 17 or applying Proposition 17 itself, we get the Salem-Zygmund condition.

For (ii) \Rightarrow (i), the idea is that ii) is equivalent to the condition

$$\int_0^1 \frac{\Upsilon(t)}{t\sqrt{\ln(1/t)}} dt < +\infty,$$

which in turn is equivalent to the entropy condition on the whole manifold \mathcal{M} thanks to Proposition 6 (note that the Riemannian probability volume μ of \mathcal{M} satisfies (28)). We then get (i) thanks to the Dudley theorem.

– For the proof of the analogue of Proposition 18, the argument is much simpler since Theorem 10 gives an equivalence holding true on the whole manifold \mathcal{M} .

– Finally, the quantitative counterpart given by Theorem 3 can be proved completely similarly once we use the inequalities given by the Slepian theorem (see [LT91, Cor.3.14] or [LQ18b, p. 12]).

11. PROOF OF (18) IN THEOREM 1, QUANTITATIVE VERSION

In the following proof, the implicit constants merely depend on $\mathcal{M}, \mathcal{M}_s$ and K (except for the setting $\mathcal{M} = S^d$ and $f_n \in \ker(\Delta + n(n + d - 1))$ for which K is obviously irrelevant). Moreover, in the manifold framework, the spectral parameter K is assumed to be large enough so that the conclusions of Theorem 2 and Theorem 3 hold.

The quantitative part of Theorem 1, namely the equivalence (18), is a direct consequence of Proposition 19, Proposition 22 below and the Cauchy-Schwarz inequality:

$$(79) \quad \mathbf{E}_\omega \left[\sup_{x \in \mathcal{M}_s} \left| \sum_{n=1}^N X_n(\omega) f_n^\omega(x) \right| \right] \hookrightarrow \mathbf{E}_\omega \left[\sup_{x \in \mathcal{M}_s} \left| \sum_{n=1}^N X_n(\omega) f_n^\omega(x) \right|^2 \right]^{1/2}.$$

PROPOSITION 19. — *There is a constant $C > 0$ such that for any $N \in \mathbb{N}^*$, any sequence (f_n) as in Theorem 1 and random variables (X_n) belonging to $L^2(\Omega)$, the following holds true*

$$(80) \quad \mathbf{E}_\omega \left[\sup_{x \in \mathcal{M}_s} \left| \sum_{n=1}^N X_n(\omega) f_n^\omega(x) \right|^2 \right]^{1/2} \hookrightarrow C \sqrt{\sup_{1 \leq n \leq N} \mathbf{E}[|X_n|^2]} \times \sum_{p=1}^N \frac{1}{p\sqrt{\ln(p+1)}} \left(\sum_{n=p}^N \|f_n\|_{L^2(\mathcal{M})}^2 \right)^{1/2},$$

where all the random variables $X_1(\omega), f_1^\omega, \dots, X_N(\omega), f_N^\omega$ are assumed to be independent.

The last result is analogue to [MP81, p. 53, Lem. 1.1] or [LQ18b, p. 249, Th. III.6] but the proof of Proposition 19 is somehow simpler and relies, among other arguments, on fact that the Salem-Zygmund condition (25) is very explicit. We begin by recalling a form of the *contraction principle* that we will use for the symmetric random functions $F_n : \omega \mapsto f_n^\omega$ (see the definitions in (12) and (14)).

PROPOSITION 20. — *Let B be a Banach space and F_1, \dots, F_N be N independent and symmetric random variables belonging to $L^p(\Omega, B)$ for some $p \in [1, +\infty)$. Then for any tuple $(a_1, \dots, a_N) \in \mathbb{R}^N$, the following inequalities hold true*

$$\min_{1 \leq n \leq N} |a_n| \cdot \mathbf{E} \left[\left\| \sum_{n=1}^N F_n \right\|_B^p \right]^{1/p} \leq \mathbf{E} \left[\left\| \sum_{n=1}^N a_n F_n \right\|_B^p \right]^{1/p} \leq \max_{1 \leq n \leq N} |a_n| \cdot \mathbf{E} \left[\left\| \sum_{n=1}^N F_n \right\|_B^p \right]^{1/p}.$$

Proof. — The first inequality is a consequence of the second one (upon assuming that each a_n is not zero and upon replacing a_n with $1/a_n$). For the second one, we refer to [LQ18a, p. 136] or the proof of [LT91, Th. 4.4].

It is known that the previous result has a reformulation if each coefficient a_n is random (see [LT91, Lem. 4.5], [MP81, Th. 4.9, p. 45] or [LQ18a, p. 137, Th. IV.4]). For the convenience of the reader, we write the proof of the following result (but all ideas are included in the last references).

COROLLARY 21. — *Let B be a Banach space, F_1, \dots, F_N be N symmetric random variables belonging to $L^p(\Omega, B)$ for some $p \in [1, +\infty)$ and A_1, \dots, A_N be N real random variables belonging to $L^p(\Omega, \mathbb{R})$. We assume that the $2N$ variables $A_1, F_1, \dots, A_N, F_N$ are mutually independent. Then the following inequalities hold true*

$$(81) \quad \mathbf{E} \left[\left\| \sum_{n=1}^N A_n F_n \right\|_B^p \right]^{1/p} \leq \mathbf{E} \left[\max_{1 \leq n \leq N} |A_n|^p \right]^{1/p} \cdot \mathbf{E} \left[\left\| \sum_{n=1}^N F_n \right\|_B^p \right]^{1/p},$$

$$(82) \quad \min_{1 \leq n \leq N} \mathbf{E}[|A_n|] \cdot \mathbf{E} \left[\left\| \sum_{n=1}^N F_n \right\|_B^p \right]^{1/p} \leq \mathbf{E} \left[\left\| \sum_{n=1}^N A_n F_n \right\|_B^p \right]^{1/p}.$$

Proof. — By independence (see [Ime19, App. F] for more details), we may write

$$(83) \quad \mathbf{E} \left[\left\| \sum_{n=1}^N A_n F_n \right\|_B^p \right] = \mathbf{E}_{\omega_1} \mathbf{E}_{\omega_2} \left[\left\| \sum_{n=1}^N A_n(\omega_1) F_n(\omega_2) \right\|_B^p \right].$$

We now freeze ω_1 and apply Proposition 20 with respect to the expectation in ω_2 . Hence we get

$$\mathbf{E}_{\omega_2} \left[\left\| \sum_{n=1}^N A_n(\omega_1) F_n(\omega_2) \right\|_B^p \right] \leq \max_{1 \leq n \leq N} |A_n(\omega_1)|^p \times \mathbf{E}_{\omega_2} \left[\left\| \sum_{n=1}^N F_n(\omega_2) \right\|_B^p \right].$$

We then obtain the expected inequality by integrating in ω_1 .

In order to get the lower bound, we first need to modify the right-hand side of (83). For ω_1 fixed, let us consider $\theta_n(\omega_1) = \pm 1$ such that $A_n(\omega_1)\theta_n(\omega_1) = |A_n(\omega_1)|$.

By symmetry and independence, one can replace each $F_n(\omega_2)$ with $\theta_n(\omega_1)F_n(\omega_2)$ in the expectation \mathbf{E}_{ω_2} in (83). So we get

$$(84) \quad \mathbf{E} \left[\left\| \sum_{n=1}^N A_n F_n \right\|_B^p \right] = \mathbf{E}_{\omega_1} \mathbf{E}_{\omega_2} \left[\left\| \sum_{n=1}^N |A_n(\omega_1)| F_n(\omega_2) \right\|_B^p \right].$$

We now bound from below with the Hölder inequality and the triangular inequality in ω_1 , namely $\mathbf{E}_{\omega_1}[\|\cdot\|] > \|\mathbf{E}_{\omega_1}[\cdot]\|$, as follows

$$\begin{aligned} \mathbf{E} \left[\left\| \sum_{n=1}^N A_n F_n \right\|_B^p \right] &= \mathbf{E}_{\omega_2} \left[\mathbf{E}_{\omega_1} \left[\left\| \sum_{n=1}^N |A_n(\omega_1)| F_n(\omega_2) \right\|_B^p \right] \right] \\ &> \mathbf{E}_{\omega_2} \left[\mathbf{E}_{\omega_1} \left[\left\| \sum_{n=1}^N |A_n(\omega_1)| F_n(\omega_2) \right\|_B \right]^p \right] \\ &> \mathbf{E}_{\omega_2} \left[\left\| \sum_{n=1}^N \mathbf{E}_{\omega_1} [|A_n(\omega_1)| F_n(\omega_2)] \right\|_B^p \right] = \mathbf{E} \left[\left\| \sum_{n=1}^N \mathbf{E} [|A_n|] F_n \right\|_B^p \right]. \end{aligned}$$

The second inequality of the statement is then a consequence of Proposition 20.

Proof of Proposition 19

Step 1. — We first need to compare f_n^ω with $f_n^{G,\omega}$. More precisely, we shall prove

$$(85) \quad \mathbf{E}_\omega \left[\sup_{x \in \mathcal{M}_s} \left| \sum_{n=1}^N X_n(\omega) f_n^\omega(x) \right|^2 \right] \leq C \mathbf{E}_{\omega'} \mathbf{E}_\omega \left[\sup_{x \in \mathcal{M}_s} \left| \sum_{n=1}^N |X_n(\omega')| f_n^{G,\omega}(x) \right|^2 \right].$$

The distribution equivalence (22) for the manifold framework suggests introducing the following random variable (collinear to a so-called chi random variable)

$$(86) \quad \forall \omega \in \Omega, \quad \chi_n(\omega) := \frac{1}{\sqrt{\dim(E_{(Kn-K,Kn)})}} \left(\sum_{Kn-K < \lambda_k \leq Kn} g_{n,k}^2(\omega) \right)^{1/2}.$$

The case $\mathcal{M} = S^d$ with $f_n \in \ker(\Delta + n(n+d-1))$ is completely similar and merely needs to change the definition (86). Having assumed the independence of all the random variables, we may write

$$(87) \quad \begin{aligned} \mathbf{E}_{\omega'} \mathbf{E}_\omega \left[\sup_{x \in \mathcal{M}_s} \left| \sum_{n=1}^N |X_n(\omega')| f_n^{G,\omega}(x) \right|^2 \right] \\ = \mathbf{E}_{\omega'} \mathbf{E}_\omega \left[\sup_{x \in \mathcal{M}_s} \left| \sum_{n=1}^N |X_n(\omega')| \chi_n(\omega) f_n^\omega(x) \right|^2 \right]. \end{aligned}$$

For almost any $\omega' \in \Omega$, one may see $|X_n(\omega')|$ as a constant in the expectation \mathbf{E}_ω . Hence, the contraction principle given by (82) in $L^2(\Omega, \mathcal{C}^0(\mathcal{M}_s))$, with $A_n(\omega) = \chi_n(\omega)$ and $F_n(\omega) = |X_n(\omega')| f_n^\omega(x)$, gives the bound from below

$$\begin{aligned} \mathbf{E}_{\omega'} \mathbf{E}_\omega \left[\sup_{x \in \mathcal{M}_s} \left| \sum_{n=1}^N |X_n(\omega')| \chi_n(\omega) f_n^\omega(x) \right|^2 \right] \\ > \left(\inf_{1 \leq n \leq N} \mathbf{E}[\chi_n^2] \right) \mathbf{E}_{\omega'} \mathbf{E}_\omega \left[\sup_{x \in \mathcal{M}_s} \left| \sum_{n=1}^N |X_n(\omega')| f_n^\omega(x) \right|^2 \right]. \end{aligned}$$

By independence and symmetry of each $\omega \mapsto f_n^\omega$ (as in (84)), the last lower bound equals

$$\left(\inf_{1 \leq n \leq N} \mathbf{E}[X_n]^2 \right) \mathbf{E}_\omega \left[\sup_{x \in \mathcal{M}_s} \left| \sum_{n=1}^N X_n(\omega) f_n^\omega(x) \right|^2 \right].$$

Now we use the inequality $\mathbf{E}[X_n] > \sqrt{2}/\sqrt{\pi}$ (see Appendix A) to get (85).

Step 2. — For ω' fixed, we now recall that $\sum_{n=1}^N |X_n(\omega')| f_n^{G,\omega}$ can be seen, with respect to ω , as a Gaussian random finite sum in the Banach space $\mathcal{C}^0(\mathcal{M}_s)$. We now invoke the important property stating that all the moments of such a Gaussian sum are universally equivalent, that is the Gaussian version of the Kahane-Khintchine inequalities (see [LT91, Cor. 3.2], [MP81, p. 44] or [LQ18a, p. 256, Cor. V.27]). Hence, (85) implies the following bound

$$\mathbf{E}_\omega \left[\sup_{x \in \mathcal{M}_s} \left| \sum_{n=1}^N X_n(\omega) f_n^\omega(x) \right|^2 \right] \leq C \mathbf{E}_{\omega'} \left[\mathbf{E}_\omega \left[\sup_{x \in \mathcal{M}_s} \left| \sum_{n=1}^N |X_n(\omega')| f_n^{G,\omega}(x) \right|^2 \right] \right].$$

Note now that changing f_n by $X_n(\omega') f_n$ in (21) leads to change $f_n^{G,\omega}$ by $|X_n(\omega')| f_n^{G,\omega}$. Consequently, Theorem 3 (for f_n replaced with $X_n(\omega') f_n$) allows us to replace the last upper bound with

$$(88) \quad C \mathbf{E}_{\omega'} \left[\left(\sum_{p=1}^N \frac{1}{p \sqrt{\ln(p+1)}} \left(\sum_{n=p}^N X_n(\omega')^2 \|f_n\|_{L^2(\mathcal{M})}^2 \right)^{1/2} \right)^2 \right].$$

Step 3 below will show the concavity of the following function:

$$(89) \quad \Psi : [0, +\infty)^N \longrightarrow \mathbb{R} \\ (t_1, \dots, t_N) \longmapsto \left(\sum_{p=1}^N \frac{1}{p \sqrt{\ln(p+1)}} \left(\sum_{n=p}^N t_n \|f_n\|_{L^2(\mathcal{M})}^2 \right)^{1/2} \right)^2.$$

Note that the number in (88) equals $C \mathbf{E}[\Psi(X_1^2, \dots, X_N^2)]$. Hence, the multidimensional Jensen inequality allows to bound it by

$$C \Psi(\mathbf{E}[X_1^2], \dots, \mathbf{E}[X_N^2]) = C \left(\sum_{p=1}^N \frac{1}{p \sqrt{\ln(p+1)}} \left(\sum_{n=p}^N \mathbf{E}[|X_n|^2] \|f_n\|_{L^2(\mathcal{M})}^2 \right)^{1/2} \right)^2 \\ \leq C \left(\sup_{1 \leq n \leq N} \mathbf{E}[|X_n|^2] \right) \left(\sum_{p=1}^N \frac{1}{p \sqrt{\ln(p+1)}} \left(\sum_{n=p}^N \|f_n\|_{L^2(\mathcal{M})}^2 \right)^{1/2} \right)^2.$$

The expected inequality (80) is proved.

Step 3. — It remains to check the concavity of (89). By developing the square in (89), we see that it is sufficient to prove the concavity of any continuous function of the form $\sqrt{\Phi_1 \Phi_2}$ on the set $\Lambda := \{\Phi_1 > 0\} \cap \{\Phi_2 > 0\}$ in which $\Phi_1 : \mathbb{R}^N \rightarrow \mathbb{R}$ and $\Phi_2 : \mathbb{R}^N \rightarrow \mathbb{R}$ are two linear functionals. Actually, such a fact is a straightforward consequence

of the Cauchy-Schwarz inequality applied to the two vectors $(\sqrt{\Phi_1(s)}, \sqrt{\Phi_1(t)})$ and $(\sqrt{\Phi_2(s)}, \sqrt{\Phi_2(t)})$ for any $(s, t) \in \Lambda^2$:

$$\begin{aligned} \sqrt{\Phi_1((s+t)/2)\Phi_2((s+t)/2)} &= \frac{\sqrt{\Phi_1(s) + \Phi_1(t)}\sqrt{\Phi_2(s) + \Phi_2(t)}}{2} \\ &> \frac{\sqrt{\Phi_1(s)}\sqrt{\Phi_2(s)} + \sqrt{\Phi_1(t)}\sqrt{\Phi_2(t)}}{2}. \end{aligned}$$

In order to finish the proof of the numerical equivalence (18), we have to reverse the inequality (80). Actually, we will use a suitable truncation argument first used by Marcus and Pisier (see [MP81, p. 53-54, proof of Lem. 1.1 & p. 99, Lem. 3.7]). In our specific context, the truncation argument is based on the uniform estimate (91) of the sequence of random variables $(\chi_n)_{n>1}$ defined in (86).

PROPOSITION 22. — *There is a constant $C > 0$ such that for any $N \in \mathbb{N}^*$ and any sequence (f_n) and random variables (X_n) as in Theorem 1, the following holds true*

$$\begin{aligned} \mathbf{E}_\omega \left[\sup_{x \in \mathcal{M}_s} \left| \sum_{n=1}^N X_n(\omega) f_n^\omega(x) \right| \right] \\ > C \left(\inf_{1 \leq n \leq N} \mathbf{E}[|X_n|] \right) \times \sum_{p=1}^N \frac{1}{p\sqrt{\ln(p+1)}} \left(\sum_{n=p}^N \|f_n\|_{L^2(\mathcal{M})}^2 \right)^{1/2}. \end{aligned}$$

Proof. — The contraction principle (82) and Theorem 3 ensure that we merely have to prove the following inequality:

$$(90) \quad \mathbf{E}_\omega \left[\sup_{x \in \mathcal{M}_s} \left| \sum_{n=1}^N f_n^{G,\omega}(x) \right| \right] \leq C \mathbf{E}_\omega \left[\sup_{x \in \mathcal{M}_s} \left| \sum_{n=1}^N f_n^\omega(x) \right| \right].$$

As in (87), we have

$$\mathbf{E}_\omega \left[\sup_{x \in \mathcal{M}_s} \left| \sum_{n=1}^N f_n^{G,\omega}(x) \right| \right] = \mathbf{E}_\omega \left[\sup_{x \in \mathcal{M}_s} \left| \sum_{n=1}^N \chi_n(\omega) f_n^\omega(x) \right| \right],$$

where all the random variables are assumed to be mutually independent. Let us now consider a number $M > 0$ (that will be chosen below) and we bound

$$\begin{aligned} \mathbf{E}_\omega \left[\sup_{x \in \mathcal{M}_s} \left| \sum_{n=1}^N f_n^{G,\omega}(x) \right| \right] &\leq \mathbf{E}_\omega \left[\sup_{x \in \mathcal{M}_s} \left| \sum_{n=1}^N \chi_n(\omega) \mathbf{1}_{\chi_n(\omega) > M} f_n^\omega(x) \right| \right] \\ &\quad + \mathbf{E}_\omega \left[\sup_{x \in \mathcal{M}_s} \left| \sum_{n=1}^N \chi_n(\omega) \mathbf{1}_{\chi_n(\omega) \leq M} f_n^\omega(x) \right| \right]. \end{aligned}$$

By using (79) and (80) with $X_n = \chi_n \mathbf{1}_{\chi_n > M}$ and Theorem 3 on the one hand, and then the contraction principle (81) on the other hand, we obtain

$$\begin{aligned} \mathbf{E}_\omega \left[\sup_{x \in \mathcal{M}_s} \left| \sum_{n=1}^N f_n^{G,\omega}(x) \right| \right] \\ \leq C \left(\sup_{n \in \mathbb{N}^*} \sqrt{\mathbf{E}[|\chi_n \mathbf{1}_{\chi_n > M}|^2]} \right) \mathbf{E}_\omega \left[\sup_{x \in \mathcal{M}_s} \left| \sum_{n=1}^N f_n^{G,\omega}(x) \right| \right] + M \mathbf{E}_\omega \left[\sup_{x \in \mathcal{M}_s} \left| \sum_{n=1}^N f_n^\omega(x) \right| \right]. \end{aligned}$$

In order to get (90), it is sufficient to justify that there is $M > 0$ satisfying

$$\sup_{n \in \mathbb{N}^*} \sqrt{\mathbf{E}[|\chi_n \mathbf{1}_{\chi_n > M}|^2]} < \frac{1}{C}.$$

That is actually a consequence of the following more precise bound (See Appendix A):

$$(91) \quad \mathbf{E}[|\chi_n \mathbf{1}_{\chi_n > M}|^2] = \mathbf{E}[\chi_n^2 \mathbf{1}_{\chi_n > M}] \leq \mathbf{E}[\chi_n^4 / M^2] \leq \frac{3}{M^2}.$$

12. PROOF OF THEOREM 1, QUALITATIVE VERSION

The end of the proof of Theorem 1 is a consequence of the equivalence (i) \Leftrightarrow (iv) of the following result.

PROPOSITION 23. — *Let us assume the same assumptions as in Theorem 1. Then the following assertions are equivalent:*

- (i) *the random series $\sum X_n(\omega) f_n^\omega$ almost surely converges in $\mathcal{C}^0(\mathcal{M}_s)$,*
- (ii) *with probability 1, the sequence of partial sums of the random series $\sum X_n(\omega) f_n^\omega$ is bounded in $\mathcal{C}^0(\mathcal{M}_s)$,*
- (iii) *the sequence of partial sums of the random series $\sum X_n(\omega) f_n^\omega$ is bounded in $L^1(\Omega, \mathcal{C}^0(\mathcal{M}_s))$,*
- (iv) *the Salem-Zygmund condition*

$$\sum_{p=1}^{+\infty} \frac{1}{p \sqrt{\ln(p+1)}} \left(\sum_{n=p}^{+\infty} \|f_n\|_{L^2(\mathcal{M})}^2 \right)^{1/2} < +\infty$$

holds,

- (v) *the random series $\sum X_n(\omega) f_n^\omega$ converges in $L^2(\Omega, \mathcal{C}^0(\mathcal{M}_s))$.*

REMARK 24. — The Gaussian random series $\sum f_n^{G,\omega}$ can be written as $\sum \chi_n(\omega) f_n^\omega$ (see (87)) in which the sequence $(X_n) = (\chi_n)$ satisfies the assumptions (17) (see Appendix A). In other words, Proposition 23 holds true for the Gaussian random series $\sum f_n^{G,\omega}$.

Let us prove Proposition 23.

(i) \Rightarrow (ii). — Obvious.

(ii) \Rightarrow (iii). — We shall use the Paley-Zygmund inequality in a similar spirit to [MP81, p. 55, Lem. 1.2] and [Ime18, Prop. 2.17]. For any $N \in \mathbb{N}^*$ and $\omega \in \Omega$, we set the partial sum

$$S_N(\omega) := \sum_{n=1}^N X_n(\omega) f_n^\omega.$$

We now give a proof by contradiction and we may assume that there is a subsequence of integers $(N_k)_{k>1}$ satisfying

$$(92) \quad \mathbf{E}[\|S_{N_k}\|_{\mathcal{C}^0(\mathcal{M}_s)}] > 2k.$$

The classical Paley-Zygmund inequality [Kah85, p. 8, Ineq. II] can be combined to Proposition 19 and Proposition 22 so we get

$$\mathbf{P}\left(\|S_{N_k}\|_{\mathcal{C}^0(\mathcal{M}_s)} > \frac{1}{2}\mathbf{E}[\|S_{N_k}\|_{\mathcal{C}^0(\mathcal{M}_s)}]\right) > \frac{\mathbf{E}[\|S_{N_k}\|_{\mathcal{C}^0(\mathcal{M}_s)}]^2}{4\mathbf{E}[\|S_{N_k}\|_{\mathcal{C}^0(\mathcal{M}_s)}^2]} > C \frac{\inf_{1 \leq n \leq N_k} \mathbf{E}[|X_n|^2]}{\sup_{1 \leq n \leq N_k} \mathbf{E}[|X_n|^2]}.$$

Therefore (17) and (92) imply

$$\inf_{k > 1} \mathbf{P}(\|S_{N_k}\|_{\mathcal{C}^0(\mathcal{M}_s)} > k) > 0.$$

That inequality contradicts ii) since the dominated convergence theorem proves

$$\lim_{k \rightarrow +\infty} \mathbf{P}(\|S_{N_k}\|_{\mathcal{C}^0(\mathcal{M}_s)} > k) = \lim_{k \rightarrow +\infty} \mathbf{E}_\omega \left[\mathbf{1}_{\{\|S_{N_k}(\omega)\|_{\mathcal{C}^0(\mathcal{M}_s)} > k\}} \right] = 0.$$

(iii) \Rightarrow (iv). — Thanks to Proposition 22 and the assumption (17), we have

$$\sup_{N \in \mathbb{N}^*} \sum_{p=1}^N \frac{1}{p\sqrt{\ln(p+1)}} \left(\sum_{n=p}^N \|f_n\|_{L^2(\mathcal{M})}^2 \right)^{1/2} < +\infty,$$

which implies

$$\sup_{\substack{(N, N') \in (\mathbb{N}^*)^2 \\ N < N'}} \sum_{p=1}^N \frac{1}{p\sqrt{\ln(p+1)}} \left(\sum_{n=p}^{N'} \|f_n\|_{L^2(\mathcal{M})}^2 \right)^{1/2} < +\infty,$$

and which in turn means the convergence of

$$\sum_{p=1}^{+\infty} \frac{1}{p\sqrt{\ln(p+1)}} \left(\sum_{n=p}^{+\infty} \|f_n\|_{L^2(\mathcal{M})}^2 \right)^{1/2}.$$

(iv) \Rightarrow (v). — Let us prove that $(S_N)_{N > 1}$ is a Cauchy sequence in $L^2(\Omega, \mathcal{C}^0(\mathcal{M}_s))$. As a consequence of Proposition 19, for any integers $N' > N$ we have

$$\begin{aligned} & \mathbf{E} \left[\|S_{N'} - S_N\|_{\mathcal{C}^0(\mathcal{M}_s)}^2 \right]^{1/2} \\ & \leq C \sqrt{\sup_{N < n \leq N'} \mathbf{E}[|X_n|^2]} \times \sum_{p=1}^{N'} \frac{1}{p\sqrt{\ln(p+1)}} \left(\sum_{n=p}^{N'} \|f_n\|_{L^2(\mathcal{M})}^2 \mathbf{1}_{[N+1, +\infty)}(n) \right)^{1/2}. \end{aligned}$$

Looking at $p = 1$ in the Salem-Zygmund condition, we see that (iv) implies the convergence of the series $\sum \|f_n\|_{L^2(\mathcal{M})}^2$. We then conclude with the assumption (17) and the following limit proved via the discrete dominated convergence theorem:

$$\lim_{N \rightarrow +\infty} \sum_{p=1}^{+\infty} \frac{1}{p\sqrt{\ln(p+1)}} \left(\sum_{n=p}^{+\infty} \|f_n\|_{L^2(\mathcal{M})}^2 \mathbf{1}_{[N+1, +\infty)}(n) \right)^{1/2} = 0.$$

(v) \Rightarrow (i). — Due to the Markov inequality, we get the convergence in probability in $\mathcal{C}^0(\mathcal{M}_s)$. Then (i) is a known consequence (see [LT91, Th. 6.1] or [LQ18a, p. 130–131, Th. III.3]).

13. PROOF OF THEOREM 5, SEMI-CLASSICAL ANALYSIS AND INDEPENDENCE OF THE RIEMANNIAN METRIC

The Salem-Zygmund condition (19) depends on the Riemannian structure via the spectral analysis of the Laplace-Beltrami operator Δ . Hence, we have to prove that if one considers another Laplace-Beltrami operator Δ on \mathcal{M} (defined with another Riemannian structure of \mathcal{M}), then the Salem-Zygmund condition does not change.

For any $K > K_0$ we set $f = \sum_{n>1} f_n \in L^2(\mathcal{M})$ with $f_n \in E_{(K_n-K, K_n]}$. Noting that f_1 does not impact the convergence of the Salem-Zygmund condition (19) and remembering that the subspaces $E_{(K_n-K, K_n]}$ are orthogonal, we see that (19) is indeed equivalent to

$$(93) \quad \sum_{p=1}^{+\infty} \frac{\|\Pi_{(Kp, +\infty)}(f)\|_{L^2(\mathcal{M})}}{p\sqrt{\ln(p+1)}} < +\infty,$$

where $\Pi_{(Kp, +\infty)} : L^2(\mathcal{M}) \rightarrow L^2(\mathcal{M})$ is the spectral projector on the spectral window $(Kp, +\infty)$ with respect to $\sqrt{-\Delta}$. Although the inequality $K > K_0$ is essential in the proof of Theorem 1 (see Theorem 10), it turns out that (93) is independent of $K > 0$ from a Hilbertian point of view. More precisely, the following lemma shows the first part of the statement of Theorem 5.

LEMMA 25. — *For any $f \in L^2(\mathcal{M})$, the condition (93) is independent of $K \in (0, +\infty)$.*

Proof. — Let K' and K be two spectral parameters satisfying $K \circlearrowleft K'$. Let us moreover consider a positive integer q satisfying $K \circlearrowleft K' \circlearrowleft Kq$. Thus we clearly have

$$\sum_{p=1}^{+\infty} \frac{\|\Pi_{(Kp, +\infty)}(f)\|_{L^2(\mathcal{M})}}{p\sqrt{\ln(p+1)}} > \sum_{p=1}^{+\infty} \frac{\|\Pi_{(K'p, +\infty)}(f)\|_{L^2(\mathcal{M})}}{p\sqrt{\ln(p+1)}}.$$

In order to reverse that condition, we just note that for any $r \in \{0, 1, \dots, q-1\}$ we have $K'p \circlearrowleft Kpq + Kr$ and hence for $p > 1$

$$\frac{\|\Pi_{(K'p, +\infty)}(f)\|_{L^2(\mathcal{M})}}{p\sqrt{\ln(p+1)}} > \frac{\|\Pi_{(Kpq+Kr, +\infty)}(f)\|_{L^2(\mathcal{M})}}{p\sqrt{\ln(p+1)}} > \frac{\|\Pi_{(Kpq+Kr, +\infty)}(f)\|_{L^2(\mathcal{M})}}{(pq+r)\sqrt{\ln(pq+r+1)}}.$$

By summing over r and p , we get

$$\begin{aligned} \sum_{p=1}^{+\infty} \frac{\|\Pi_{(K'p, +\infty)}(f)\|_{L^2(\mathcal{M})}}{p\sqrt{\ln(p+1)}} &= \frac{1}{q} \sum_{r=0}^{q-1} \sum_{p=1}^{+\infty} \frac{\|\Pi_{(K'p, +\infty)}(f)\|_{L^2(\mathcal{M})}}{p\sqrt{\ln(p+1)}} \\ &> \frac{1}{q} \sum_{r=0}^{q-1} \sum_{p=1}^{+\infty} \frac{\|\Pi_{(Kpq+Kr, +\infty)}(f)\|_{L^2(\mathcal{M})}}{(pq+r)\sqrt{\ln(pq+r+1)}} = \frac{1}{q} \sum_{p=q}^{+\infty} \frac{\|\Pi_{(Kp, +\infty)}(f)\|_{L^2(\mathcal{M})}}{p\sqrt{\ln(p+1)}}. \end{aligned}$$

The sequel is devoted to the proof of the invariance of (93) with respect to the choice of the Laplace-Beltrami operator Δ . We first need a semi-classical reformulation of the Salem-Zygmund condition.

LEMMA 26. — *Let us fix a smooth function $\Psi : [0, +\infty) \rightarrow [0, 1]$ satisfying*

$$(94) \quad \begin{aligned} t \in [0, 1] &\implies \Psi(t) = 1, \\ t > 2 &\implies \Psi(t) = 0. \end{aligned}$$

Then, for any $f \in L^2(\mathcal{M})$, the Salem-Zygmund condition (93) is equivalent to the following semi-classical condition

$$(95) \quad \int_0^1 \frac{\|f - \Psi(-h^2 \Delta)f\|_{L^2(\mathcal{M})}}{h\sqrt{-\ln(h)}} dh < +\infty.$$

Proof. — Following (9), we decompose $f = \sum_{k \in \mathbb{N}} c_k \phi_k$ with $(c_k) \in \ell^2(\mathbb{N})$. Hence we get

$$\begin{aligned} f - \Psi(-h^2 \Delta)f &= \sum_{k=0}^{+\infty} (1 - \Psi(h^2 \lambda_k^2)) c_k \phi_k, \\ \|f - \Psi(-h^2 \Delta)f\|_{L^2(\mathcal{M})}^2 &= \sum_{k=0}^{+\infty} (1 - \Psi(h^2 \lambda_k^2))^2 |c_k|^2. \end{aligned}$$

The imposed conditions on Ψ allow us to bound as follows

$$\begin{aligned} \sum_{\lambda_k > \sqrt{2}/h} |c_k|^2 &\subset \|f - \Psi(-h^2 \Delta)f\|_{L^2(\mathcal{M})}^2 \subset \sum_{\lambda_k > 1/h} |c_k|^2, \\ \|\Pi_{(\sqrt{2}/h, +\infty)}(f)\|_{L^2(\mathcal{M})} &\subset \|f - \Psi(-h^2 \Delta)f\|_{L^2(\mathcal{M})} \subset \|\Pi_{(1/h, +\infty)}(f)\|_{L^2(\mathcal{M})}. \end{aligned}$$

By integrating with respect to $h \in (0, 1]$, we get

$$(96) \quad \int_0^1 \frac{\|\Pi_{(\sqrt{2}/h, +\infty)}(f)\|_{L^2(\mathcal{M})}}{h\sqrt{-\ln(h)}} dh \subset \int_0^1 \frac{\|f - \Psi(-h^2 \Delta)f\|_{L^2(\mathcal{M})}}{h\sqrt{-\ln(h)}} dh$$

$$(97) \quad \subset \int_0^1 \frac{\|\Pi_{(1/h, +\infty)}(f)\|_{L^2(\mathcal{M})}}{h\sqrt{-\ln(h)}} dh.$$

By using that $h \in (0, 1] \mapsto \|\Pi_{(1/h, +\infty)}(f)\|_{L^2(\mathcal{M})}$ is non-decreasing, we see that the upper bound in (97) is bounded from above by

$$\begin{aligned} \sum_{p>1} \int_{1/(p+1)}^{1/p} \frac{\|\Pi_{(1/h, +\infty)}(f)\|_{L^2(\mathcal{M})}}{h\sqrt{-\ln(h)}} dh &\subset \sum_{p>1} \|\Pi_{(p, +\infty)}(f)\|_{L^2(\mathcal{M})} \int_{1/(p+1)}^{1/p} \frac{dh}{h\sqrt{-\ln(h)}} \\ &\quad \cdot \sum_{p>1} \frac{\|\Pi_{(p, +\infty)}(f)\|_{L^2(\mathcal{M})}}{p\sqrt{\ln(p+1)}}. \end{aligned}$$

Similarly, we see that the lower bound in (96) is bounded from below by

$$\sum_{p>1} \int_{1/(p+1)}^{1/p} \frac{\|\Pi_{(\sqrt{2}/h, +\infty)}(f)\|_{L^2(\mathcal{M})}}{h\sqrt{-\ln(h)}} dh \ \& \ \sum_{p>2} \frac{\|\Pi_{(\sqrt{2}p, +\infty)}(f)\|_{L^2(\mathcal{M})}}{p\sqrt{\ln(p+1)}}.$$

We conclude with Lemma 25.

Before analyzing the semi-classical Salem-Zygmund condition (95), we need to recall a few facts about semi-classical symbols and semi-classical pseudo-differential operators. The standard relation between a symbol $s : \mathbb{R}^d \times \mathbb{R}^d \times (0, 1] \rightarrow \mathbb{R}$ and its quantized operator, usually denoted by $s(x, D, h)$, is given by the formula

$$(98) \quad \forall F \in \mathcal{S}(\mathbb{R}^d), \forall x \in \mathbb{R}^d, \quad (s(x, D, h)F)(x) = \int_{\mathbb{R}^d} e^{i\langle x, \xi \rangle} s(x, \xi, h) \widehat{F}(\xi) \frac{d\xi}{(2\pi)^d}.$$

For the particular case $s(x, h\xi)$, one prefers writing $s(x, hD)$. Such operators, with h running over $(0, 1]$, are called semi-classical pseudo-differential operators. Here is the result we need in the sequel.

LEMMA 27. — *For any symbols s_1 and s_2 belonging to $\mathcal{C}_b^\infty(\mathbb{R}^d \times \mathbb{R}^d)$ and having disjoint supports, for any $N \in \mathbb{N}^*$, there are constant $C > 0$ and C_N such that the following estimates uniformly hold for any $h \in (0, 1]$:*

$$(99) \quad \|s_1(x, hD)\|_{L^2(\mathbb{R}^d) \rightarrow L^2(\mathbb{R}^d)} \leq C,$$

$$(100) \quad \|s_1(x, hD) \circ s_2(x, hD)\|_{L^2(\mathbb{R}^d) \rightarrow L^2(\mathbb{R}^d)} \leq C_N h^N.$$

Proof. — We will use the notations of [Ler10, p. 22–23]. For any $m \in \mathbb{R}$, one denotes by S_{scl}^m the space of symbols $s(x, \xi, h)$ satisfying for any $\alpha \in \mathbb{N}^d$ and $\beta \in \mathbb{N}^d$:

$$\sup_{x \in \mathbb{R}^d} \sup_{\xi \in \mathbb{R}^d} \sup_{h \in (0, 1]} |(\partial_\xi^\alpha \partial_x^\beta s)(x, \xi, h)| h^{m-|\alpha|} < +\infty.$$

With those notations, the symbols $s_1(x, h\xi)$ and $s_2(x, h\xi)$ belong to the class S_{scl}^0 . The first estimate is merely the semi-classical Calderon-Vaillancourt (see [Ler10, Th. 1.1.30]). For the second estimate, [Ler10, Th. 1.1.31] states that there is a “composition” symbol $s(x, \xi, h)$ belonging to the class S_{scl}^0 satisfying

$$s_1(x, hD) \circ s_2(x, hD) = s(x, D, h).$$

Then we use the semi-classical symbolic calculus at rank N of $s_1(x, h\xi) \in S_{\text{scl}}^0$ and $s_2(x, h\xi) \in S_{\text{scl}}^0$ (see [Ler10, Th. 1.1.32]) that ensures that the “composition” symbol $s(x, \xi, h)$ satisfies

$$s(x, \xi, h) - \sum_{|\alpha| < N} \frac{1}{\alpha! i^\alpha} \partial_\xi^\alpha \{s_1(x, h\xi)\} \partial_x^\alpha \{s_2(x, h\xi)\} \in S_{\text{scl}}^{-N},$$

which means that $s(x, \xi, h)$ belongs to S_{scl}^{-N} because the partial sum vanishes thanks to the assumption of disjoint supports. Finally, a new use of the semi-classical Calderon-Vaillancourt (see [Ler10, Th. 1.1.30]) proves (100).

We now recall the local expression of the Laplace-Beltrami operator Δ of \mathcal{M} . Let $\tau : U \subset \mathbb{R}^d \rightarrow V \subset \mathcal{M}$ be a coordinate patch of \mathcal{M} . For any $x \in U$, we denote by g_x the $d \times d$ positive-definite matrix corresponding to the Riemannian metric at $\tau(x) \in \mathcal{M}$ via the coordinate patch τ . Moreover, $(g_x^{ij})_{i,j}$ is the usual notation for the inverse of the matrix g_x . For any smooth function $f : \mathcal{M} \rightarrow \mathbb{C}$ with compact support in V ,

the action of the Laplace-Beltrami operator on f can be seen as that of a differential operator acting on $f \circ \tau : U \rightarrow \mathbb{R}$ as follows (see for instance [Shu01, p. 167]):

$$\begin{aligned} (\Delta f)(\tau(x)) &= \frac{1}{\sqrt{\det g_x}} \sum_{i=1}^d \sum_{j=1}^d \partial_{x_i} \left(g_x^{ij} \sqrt{\det g_x} \partial_{x_j} (f \circ \tau) \right) \\ &= \sum_{i=1}^d \sum_{j=1}^d g_x^{ij} \partial_{x_i} \partial_{x_j} (f \circ \tau) + \text{differential terms of order 1.} \end{aligned}$$

By introducing the so-called principal symbol of $-\Delta$ on the coordinate patch τ

$$(101) \quad \wp : (x, \xi) \in \mathbb{R}^d \times \mathbb{R}^d \mapsto \mathbf{1}_U(x) \sum_{i=1}^d \sum_{j=1}^d g_x^{ij} \xi_i \xi_j \in \mathbb{R}$$

and using the pseudo-differential notation (98) above (without h), the principal term of $(-\Delta f)(\tau(x))$ reads

$$-\sum_{i=1}^d \sum_{j=1}^d g_x^{ij} \partial_{x_i} \partial_{x_j} (f \circ \tau) = \wp(x, D)(f \circ \tau),$$

where $f \circ \tau$ belongs to $\mathcal{S}(\mathbb{R}^d)$ after being extended by 0 outside U . Moreover, on any compact subset of the domain U of the coordinate patch, there is by continuity a constant $\gamma > 1$ for which the following uniform estimates hold

$$(102) \quad \forall \xi \in \mathbb{R}^d, \quad \frac{1}{\gamma} |\xi|^2 \leq \wp(x, \xi) \leq \gamma |\xi|^2.$$

These inequalities implies that $-\Delta$ is a second order elliptic operator. We stress that \wp depends on the chosen coordinate patch $\tau : U \subset \mathbb{R}^d \rightarrow V \subset \mathcal{M}$ (although we merely write \wp for simplicity).

In order to avoid repeating the same technical assumptions, we set the following definition.

DEFINITION 28. — We denote by $\Lambda(\mathcal{M})$ the set of triplets $(\tau, \psi, \tilde{\psi})$ as follows

(i) $\tau : U \rightarrow V$ is a coordinate patch of \mathcal{M} from an open subset $U \subset \mathbb{R}^d$ to an open subset $V \subset \mathcal{M}$,

(ii) ψ belongs to $\mathcal{C}_c^\infty(V)$,

(iii) $\tilde{\psi}$ belongs to $\mathcal{C}_c^\infty(V)$ and equals 1 on a neighborhood of the support of ψ .

Furthermore, for any $(\tau, \psi, \tilde{\psi}) \in \Lambda(\mathcal{M})$ and any smooth function $\Psi : [0, +\infty) \rightarrow [0, 1]$ satisfying (94), we define the following two symbols

$$(103) \quad \forall (x, \xi) \in \mathbb{R}^d \times \mathbb{R}^d, \quad s_{\tau, \psi}^{\wp, \Psi}(x, \xi) := \mathbf{1}_U(x) \psi(\tau(x)) \times (1 - \Psi \circ \wp(x, \xi)),$$

$$(104) \quad s_{\tau, \psi}^{\star \Psi}(x, \xi) := \mathbf{1}_U(x) \psi(\tau(x)) \times (1 - \Psi(|\xi|^2)),$$

where \wp in (103) is given in (101) (note that $s_{\tau, \psi}^{\wp, \Psi}(x, \xi) = 0$ for $x \notin U$).

For the sequel, we warn the reader that the symbol $s_{\tau, \psi}^{\wp, \Psi}$ will appear by making a semi-classical expansion of $\Psi(-h^2 \Delta)$ in (95) (see the proof of Proposition 30). The main drawback of $s_{\tau, \psi}^{\wp, \Psi}$ is its dependence with respect to the Laplace-Beltrami

operator Δ (more precisely to \wp via the coordinate patch τ). The next result shows that the family of symbols $s_{\tau,\psi}^{\wp,\Psi}$ is equivalent, in a semi-classical sense, to the family of symbols $s_{\tau,\psi}^{\bullet\Psi}$ whose expression (104) is clearly independent of the Laplace-Beltrami operator.

PROPOSITION 29. — *Let us fix a smooth function $\Psi : [0, +\infty) \rightarrow [0, 1]$ satisfying (94). For any $(\tau, \psi, \tilde{\psi}) \in \Lambda(\mathcal{M})$, there are $\alpha > 0$ and $C > 0$ such that the following inequalities hold true uniformly in $F \in L^2(\mathbb{R}^d)$ and $h \in (0, 1]$:*

$$(105) \quad \|s_{\tau,\psi}^{\wp,\Psi}(x, hD)F\|_{L_x^2(\mathbb{R}^d)} \leq C \|s_{\tau,\tilde{\psi}}^{\bullet\Psi}(x, \alpha hD)F\|_{L_x^2(\mathbb{R}^d)} + \|F\|_{L^2(\mathbb{R}^d)} \mathcal{O}(h),$$

$$(106) \quad \|s_{\tau,\psi}^{\bullet\Psi}(x, hD)F\|_{L_x^2(\mathbb{R}^d)} \leq C \|s_{\tau,\tilde{\psi}}^{\wp,\Psi}(x, \alpha hD)F\|_{L_x^2(\mathbb{R}^d)} + \|F\|_{L^2(\mathbb{R}^d)} \mathcal{O}(h).$$

Proof. — Let us explain the main idea for (105). We will decompose

$$F = s_{\tau,\tilde{\psi}}^{\bullet\Psi}(x, \alpha hD)F + (1 - s_{\tau,\tilde{\psi}}^{\bullet\Psi}(x, \alpha hD))F,$$

which in turn will imply that $\|s_{\tau,\psi}^{\wp,\Psi}(x, hD)F\|_{L_x^2(\mathbb{R}^d)}$ is bounded by

$$\|s_{\tau,\psi}^{\wp,\Psi}(x, hD)s_{\tau,\tilde{\psi}}^{\bullet\Psi}(x, \alpha hD)F\|_{L_x^2(\mathbb{R}^d)} + \|s_{\tau,\psi}^{\wp,\Psi}(x, hD)(1 - s_{\tau,\tilde{\psi}}^{\bullet\Psi}(x, \alpha hD))F\|_{L_x^2(\mathbb{R}^d)}.$$

We will then conclude by applying (99) and (100) provided that $\alpha > 0$ is chosen large enough so that the two symbols $s_{\tau,\psi}^{\wp,\Psi}(x, \xi)$ and $1 - s_{\tau,\tilde{\psi}}^{\bullet\Psi}(x, \alpha\xi)$ have disjoint supports.

Let us now go into technical details. We first remark that the smooth function $\mathbf{1}_U(x) \times (\psi \circ \tau(x))$ has compact support and thus belongs to $\mathcal{C}_b^\infty(\mathbb{R}^d)$. Similarly, thanks to (94) and (102), one also checks that the two symbols $\mathbf{1}_U(x)(\psi \circ \tau(x)) \times \Psi(\wp(x, \xi))$ and $\mathbf{1}_U(x)(\tilde{\psi} \circ \tau(x)) \times \Psi(\alpha^2|\xi|^2)$ belong to $\mathcal{C}_b^\infty(\mathbb{R}^d \times \mathbb{R}^d)$. As a consequence of (99), the pseudo-differential operators $s_{\tau,\psi}^{\wp,\Psi}(x, hD)$ and $s_{\tau,\tilde{\psi}}^{\bullet\Psi}(x, \alpha hD)$ are well-defined and bounded on $L^2(\mathbb{R}^d)$.

We now turn to the choice of α . The two symbols $s_{\tau,\psi}^{\wp,\Psi}$ and $\mathbf{1}_U(x)(\tilde{\psi} \circ \tau(x)) \times \Psi(\alpha^2|\xi|^2)$ are respectively supported in

$$U \times \{|\xi|^2 > 1/\gamma\} \quad \text{and} \quad U \times \{|\xi|^2 \leq 2/\alpha^2\}.$$

For $\alpha > \sqrt{2\gamma}$, the two supports are disjoint. In order to prove (105), we decompose as follows for any $(x, \xi) \in \mathbb{R}^d \times \mathbb{R}^d$:

$$1 = \left(\mathbf{1}_U(x)(\tilde{\psi} \circ \tau(x)) \times (1 - \Psi(\alpha^2 h^2 |\xi|^2)) \right) + \left(\mathbf{1}_U(x)(\tilde{\psi} \circ \tau(x)) \times \Psi(\alpha^2 h^2 |\xi|^2) \right) + \left(1 - \mathbf{1}_U(x)(\tilde{\psi} \circ \tau(x)) \right),$$

which leads, after using the quantization formula (98), to

$$(107) \quad F = s_{\tau,\psi}^{\bullet\Psi}(x, \alpha hD)F + (\mathbf{1}_U \times (\tilde{\psi} \circ \tau) \times \Psi(-\alpha h^2 \Delta))F + (1 - \mathbf{1}_U \times (\tilde{\psi} \circ \tau))F.$$

Hence we get

$$(108) \quad \|s_{\tau,\psi}^{\varphi,\Psi}(x, hD)F\|_{L_x^2(\mathbb{R}^d)} \leq \|s_{\tau,\psi}^{\varphi,\Psi}(x, hD)s_{\tau,\tilde{\psi}}^{\bullet\Psi}(x, \alpha hD)F\|_{L_x^2(\mathbb{R}^d)} \\ + \|s_{\tau,\psi}^{\varphi,\Psi}(x, hD)(\mathbf{1}_U \times (\tilde{\psi} \circ \tau) \times \Psi(-\alpha h^2 \Delta))F\|_{L_x^2(\mathbb{R}^d)}$$

$$(109) \quad + \|s_{\tau,\psi}^{\varphi,\Psi}(x, hD)((1 - \mathbf{1}_U \times (\tilde{\psi} \circ \tau))F)\|_{L_x^2(\mathbb{R}^d)}.$$

By using the Calderon-Vaillancourt inequality (99) for $s_1 = s_{\tau,\psi}^{\varphi,\Psi}$, we obtain

$$\|s_{\tau,\psi}^{\varphi,\Psi}(x, hD)s_{\tau,\tilde{\psi}}^{\bullet\Psi}(x, \alpha hD)F\|_{L_x^2(\mathbb{R}^d)} \leq C \|s_{\tau,\tilde{\psi}}^{\bullet\Psi}(x, \alpha hD)F\|_{L_x^2(\mathbb{R}^d)}.$$

The inequality (105) can then be proved by bounding the two terms (108) and (109) by $\|F\|_{L^2(\mathbb{R}^d)}\mathcal{O}(h)$ thanks to (100) and the following two remarks:

(i) accordingly to the choice of $\alpha > \sqrt{2\gamma}$, the symbols $s_{\tau,\psi}^{\varphi,\Psi}(x, \xi)$ and

$$\mathbf{1}_U(x) \times (\tilde{\psi} \circ \tau(x)) \times \Psi(\alpha^2|\xi|^2)$$

have disjoint supports because of the frequency variable ξ ,

(ii) let $s_2(x, \xi)$ be the smooth symbol $1 - \mathbf{1}_U(x) \times (\tilde{\psi} \circ \tau(x))$ that merely depends on x . The quantization formula (98) shows that $s_2(x, hD)$ is the multiplication operator by $1 - \mathbf{1}_U(x) \times (\tilde{\psi} \circ \tau(x))$. We furthermore note that the spatial component of the support of $s_{\tau,\psi}^{\varphi,\Psi}(x, \xi)$ (see (103)) is included in U and more precisely in the support of $\psi \circ \tau$. But the supports of $\psi \circ \tau$ and $1 - \tilde{\psi} \circ \tau$ are disjoint because $1 - \tilde{\psi}$ vanishes in a neighborhood of the support of ψ (see Definition 28). In other words, the symbols $s_{\tau,\psi}^{\varphi,\Psi}(x, \xi)$ and $s_2(x, \xi)$ have disjoint supports.

The second inequality (106) can be proved with a similar strategy. Actually, we have to choose $\alpha > 0$ so that the two symbols $s_{\tau,\psi}^{\varphi,\Psi}(x, \xi)$ and $\mathbf{1}_U(x) \times (\tilde{\psi} \circ \tau(x)) \times (\Psi \circ \varphi)(x, \alpha \xi)$ have disjoint supports. According to (94), (102) and (104), the supports of those two symbols are respectively included in

$$U \times \{|\xi| > 1\} \quad \text{and} \quad U \times \{|\xi|^2 \leq 2\gamma/\alpha^2\}.$$

Here again, for the same choice of $\alpha > \sqrt{2\gamma}$, we may apply (99) and (100) to the following similar decomposition to (107):

$$F = s_{\tau,\psi}^{\varphi,\Psi}(x, \alpha hD)F + (\mathbf{1}_U \times \tilde{\psi} \circ \tau \times \Psi(\varphi(x, \alpha D)))F + (1 - \mathbf{1}_U \times \tilde{\psi} \circ \tau)F.$$

We are now ready to give a local but pseudo-differential reformulation of the semi-classical Salem-Zygmund condition (95).

PROPOSITION 30. — *Let us fix a smooth function $\Psi : [0, +\infty) \rightarrow [0, 1]$ satisfying (94). For any $f \in L^2(\mathcal{M})$, the semi-classical Salem-Zygmund condition (95) is equivalent to the following condition: for any $(\tau, \psi, \tilde{\psi}) \in \Lambda(\mathcal{M})$ (according to Definition 28), we have*

$$(110) \quad \int_0^1 \frac{\|s_{\tau,\psi}^{\varphi,\Psi}(x, hD)((\tilde{\psi}f) \circ \tau)\|_{L_x^2(\mathbb{R}^d)}}{h\sqrt{-\ln(h)}} dh < +\infty,$$

where $s_{\tau,\psi}^{\varphi,\Psi}$ is defined in (103).

Proof. — By compactness of \mathcal{M} , there is a finite open cover $\mathcal{M} = V_1 \cup \dots \cup V_n$ where each open subset V_i corresponds to a coordinate patch $\tau_i : U_i \subset \mathbb{R}^d \rightarrow V_i \subset \mathcal{M}$. Now consider a smooth partition of unity $1 = \psi_1 + \dots + \psi_n$ subordinate to the previous open cover of \mathcal{M} . Since the support of ψ_i is a compact subset of V_i , one infers that, for any $g \in L^2(\mathcal{M})$, the equivalence $\|\psi_i g\|_{L^2(V_i)} \simeq \|(\psi_i g) \circ \tau_i\|_{L^2(\mathbb{R}^d)}$ holds with constants independent of g . We then easily obtain the equivalence

$$\|g\|_{L^2(\mathcal{M})} \simeq \sum_{i=1}^n \|(\psi_i g) \circ \tau_i\|_{L^2(\mathbb{R}^d)}.$$

As a consequence, it is clear that (95) is equivalent to the following property: for any coordinate patch $\tau : U \subset \mathbb{R}^d \rightarrow V \subset \mathcal{M}$ and for any $\psi \in \mathcal{C}_c^\infty(V)$, we have

$$(111) \quad \int_0^1 \frac{\|(\psi \times f - \psi \times (\Psi(-h^2 \Delta) f)) \circ \tau\|_{L^2_x(\mathbb{R}^d)}}{h\sqrt{-\ln(h)}} dh < +\infty,$$

with the convention $(\psi \times f) \circ \tau(x) = 0$ for $x \notin U$. We now invoke the semi-classical functional calculus of the Laplace-Beltrami operator as stated in [BGT04, Prop. 2.1 with $N = 1$ and $\sigma = 0$]. Let $\tilde{\psi} \in \mathcal{C}_c^\infty(V)$ be a function equaling 1 on the support of ψ (in other words, $(\tau, \psi, \tilde{\psi})$ belongs to $\Lambda(\mathcal{M})$ in Definition 28). The semi-classical functional calculus ensures that there exists an explicit symbol $\Psi_0 \in \mathcal{C}_c^\infty(U \times \mathbb{R}^d)$ satisfying the following properties:

(i) the pseudo-differential $\Psi_0(x, hD)$ quantizes Ψ_0 as in (98) and satisfies the uniform estimates

$$(112) \quad \|(\psi \Psi(-h^2 \Delta) f) \circ \tau - \Psi_0(x, hD)((\tilde{\psi} f) \circ \tau)\|_{L^2(\mathbb{R}^d)} \leq Ch \|f\|_{L^2(\mathcal{M})},$$

(ii) the symbol Ψ_0 has the following explicit expression:

$$(113) \quad \forall (x, \xi) \in U \times \mathbb{R}^d, \quad \Psi_0(x, \xi) = \psi(\tau(x)) \Psi(\varrho(x, \xi)).$$

By plugging (112) in (111) and using the finiteness of $\int_0^1 dh / \sqrt{-\ln(h)} = \sqrt{\pi}$, we see that (111) is equivalent to

$$(114) \quad \int_0^1 \frac{\|(\psi f) \circ \tau - \Psi_0(x, hD)((\tilde{\psi} f) \circ \tau)\|_{L^2_x(\mathbb{R}^d)}}{h\sqrt{-\ln(h)}} dh < +\infty.$$

Remembering that $\tilde{\psi}$ equals 1 on the support of ψ , we get $\psi = \psi \tilde{\psi}$ and so $(\psi f) \circ \tau = (\psi \circ \tau) \times ((\tilde{\psi} f) \circ \tau)$. Since the pseudo-differential operator $(\mathbf{1}_U \times \psi \circ \tau)(x, hD)$ with symbol $\mathbf{1}_U \times \psi \circ \tau$ (independent of ξ) is merely the multiplication by $\mathbf{1}_U \times \psi \circ \tau$ (see (98)), one may look at (103) and (113) to conclude that (114) reduces to (110).

The following result achieves the proof of Theorem 5 because it allows us to get rid of the Laplace-Beltrami operator Δ and because the vector space $L^2(\mathcal{M})$ is independent of the Riemannian structure of \mathcal{M} .

PROPOSITION 31. — *Let us fix a smooth function $\Psi : [0, +\infty) \rightarrow [0, 1]$ satisfying (94). Then for any $f \in L^2(\mathcal{M})$, the Salem-Zygmund condition (93) is equivalent to the*

following condition: for any $(\tau, \psi, \tilde{\psi}) \in \Lambda(\mathcal{M})$ (according to Definition 28), we have

$$(115) \quad \int_0^1 \frac{\|s_{\tau, \psi}^{\bullet\Psi}(x, hD)((\tilde{\psi}f) \circ \tau)\|_{L_x^2(\mathbb{R}^d)}}{h\sqrt{-\ln(h)}} dh < +\infty,$$

where $s_{\tau, \psi}^{\bullet\Psi}$ is defined in (104).

Proof. — The key point is the following: for any $(\tau, \psi, \tilde{\psi}) \in \Lambda(\mathcal{M})$ there is $\bar{\psi} \in \mathcal{C}_c^\infty(V)$ such that $(\tau, \psi, \bar{\psi})$ and $(\tau, \bar{\psi}, \tilde{\psi})$ belong to $\Lambda(\mathcal{M})$. Indeed, there is an open subset $W \subset V$ such that $\text{supp}(\psi) \subset W \subset \{\tilde{\psi} = 1\}$. So we just have to choose $\bar{\psi} \in \mathcal{C}_c^\infty(W)$ that equals 1 on a neighborhood of the compact subset $\text{supp}(\psi)$.

If the finiteness conditions (115) hold, then for $(\tau, \bar{\psi}, \tilde{\psi}) \in \Lambda(\mathcal{M})$ we get

$$(116) \quad \int_0^1 \frac{\|s_{\tau, \bar{\psi}}^{\bullet\Psi}(x, hD)((\tilde{\psi}f) \circ \tau)\|_{L_x^2(\mathbb{R}^d)}}{h\sqrt{-\ln(h)}} dh < +\infty.$$

Thanks to the Calderon-Vaillancourt inequality (99) (as in the proof of Proposition 29), we know that $\|s_{\tau, \bar{\psi}}^{\bullet\Psi}(x, hD)((\tilde{\psi}f) \circ \tau)\|_{L_x^2(\mathbb{R}^d)}$ is bounded with respect to $h \in (0, 1]$ and hence the integrability of (116) is at $h = 0$. For any $\alpha > 0$, a simple linear change of variable shows

$$\int_0^1 \frac{\|s_{\tau, \bar{\psi}}^{\bullet\Psi}(x, \alpha hD)((\tilde{\psi}f) \circ \tau)\|_{L_x^2(\mathbb{R}^d)}}{h\sqrt{-\ln(h)}} dh < +\infty.$$

By applying (105) of Proposition 29 with $(\tau, \psi, \bar{\psi}) \in \Lambda(\mathcal{M})$, we clearly get the finiteness conditions (110) of Proposition 30. A perfectly similar argument using (106) allows us to reverse the previous implication. Finally, Lemma 26 and Proposition 30 show the equivalence with the Salem-Zygmund condition (93).

APPENDIX A. FIRST MOMENTS OF CHI DISTRIBUTIONS

We give elementary estimates about the first moments of chi distributions. For any $n \in \mathbb{N}^*$, we set

$$\chi_n := \frac{1}{\sqrt{n}} \left(\sum_{k=1}^n g_k^2 \right)^{1/2},$$

where g_1, \dots, g_n are i.i.d. Gaussian random variables $\mathcal{N}_{\mathbb{R}}(0, 1)$. The equality $\mathbf{E}[\chi_n^2] = 1$ is obvious. The following uniform inequalities also hold (with sharpness for $n = 1$):

$$\frac{\sqrt{2}}{\sqrt{\pi}} \triangleleft \mathbf{E}[\chi_n] \triangleleft 1 \quad \text{and} \quad 1 \triangleleft \mathbf{E}[\chi_n^4] \triangleleft 3.$$

– For the bound on $\mathbf{E}[\chi_n^4]$, we write

$$\mathbf{E}[\chi_n^4] = \frac{1}{n^2} \sum_{\substack{1 \leq k \leq n \\ 1 \leq \ell \leq n}} \mathbf{E}[g_k^2 g_\ell^2]$$

and use the Cauchy-Schwarz inequality $\mathbf{E}[g_k^2 g_\ell^2] \triangleleft \sqrt{\mathbf{E}[g_k^4] \mathbf{E}[g_\ell^4]} = 3$. Note that the Cauchy-Schwarz inequality also shows $1 = \mathbf{E}[\chi_n^2]^2 \triangleleft \mathbf{E}[\chi_n^4]$.

– Let us now justify the bounds of $\mathbf{E}[\chi_n]$. The inequality $\mathbf{E}[\chi_n] \leq \sqrt{\mathbf{E}[\chi_n^2]} = 1$ is direct. For the lower bound $\mathbf{E}[\chi_n] > \sqrt{2}/\sqrt{\pi}$, one considers random variables $\varepsilon_1, \dots, \varepsilon_n$ (which are, as usual, independent of all other random variables), then one has

$$\begin{aligned} \sum_{k=1}^n g_k(\omega)^2 &= \mathbf{E}_{\omega'} \left[\left(\sum_{k=1}^n \varepsilon_k(\omega') g_k(\omega) \right)^2 \right] > \mathbf{E}_{\omega'} \left[\left| \sum_{k=1}^n \varepsilon_k(\omega') g_k(\omega) \right|^2 \right] \\ \implies \mathbf{E}_{\omega} [\chi_n] &> \frac{1}{\sqrt{n}} \mathbf{E}_{\omega} \mathbf{E}_{\omega'} \left[\left| \sum_{k=1}^n \varepsilon_k(\omega') g_k(\omega) \right| \right]. \end{aligned}$$

We now remark that, for a fixed ω' , the random variable $\omega \mapsto \sum_{k=1}^n \varepsilon_k(\omega') g_k(\omega)$ is Gaussian and more precisely has the same distribution as $\sqrt{n}g_1$. Hence, we get $\mathbf{E}[\chi_n] > \mathbf{E}[|g_1|] = \sqrt{2}/\sqrt{\pi}$.

APPENDIX B. PROOF OF THEOREM 11 VIA A RESULT OF CANZANI-HANIN

As written above, Theorem 11 is considered as known (see for instance [CH18, lines (4), (5) & (8)]). We merely write elements of proofs for the non-specialist reader since we do not know a published reference. The argument here relies on the published references [CH18, Sog17] and on a Bernstein-type inequality on the boundaryless compact Riemannian manifold \mathcal{M} proved in [Bin04, line (11)].

We recall a few facts of Riemannian geometry. For any $y \in \mathcal{M}$, the Riemannian structure of \mathcal{M} induces a canonical isomorphism between the tangent space $T_y\mathcal{M}$ and the cotangent space $T_y^*\mathcal{M}$. Hence, we may endow $T_y^*\mathcal{M}$ with a canonical inner product and a canonical Euclidean norm $\xi \mapsto |\xi|_y$. Consequently, in the Riemannian setting, the exponential map of \mathcal{M} is canonically defined on the cotangent fiber bundle. We denote by $\text{inj}(M)$ the injectivity radius of \mathcal{M} , namely the supremum of radii $r > 0$ such that, for any $y \in \mathcal{M}$, the restriction of the exponential map $\exp_y : T_y^*\mathcal{M} \rightarrow M$ on the open ball $\{\xi \in T_y\mathcal{M}, |\xi|_y < r\}$ is a diffeomorphism. The injectivity radius $\text{inj}(M)$ is positive thanks to the compactness of \mathcal{M} . Moreover one has

$$\exp_y(\{\xi \in T_y^*\mathcal{M}, |\xi|_y < r\}) = B_{\delta_g}(y, r).$$

For any $\xi \in T_y^*\mathcal{M}$ in the ball $\{|\xi|_y < \text{inj}(M)\}$, the norm $|\xi|_y$ equals the distance $\delta_g(\exp_y(\xi), y)$. In particular for any point $x \in \mathcal{M}$ satisfying $\text{dist}(x, y) < \text{inj}(M)$, the element $\xi = \exp_y^{-1}(x) \in T_y^*\mathcal{M}$ is well defined and moreover satisfies the following norm equality in $T_y^*\mathcal{M}$:

$$(117) \quad |\exp_y^{-1}(x)|_y = \delta_g(x, y).$$

In order to keep the same notations as in [CH18, line (11)], we also denote by $\rho : \mathbb{R} \rightarrow \mathbb{R}$ a Schwartz function satisfying the following properties

$$(118) \quad \begin{aligned} |t| \leq \frac{1}{2} \text{inj}(M) &\implies \widehat{\rho}(t) = 1, \\ |t| > \text{inj}(M) &\implies \widehat{\rho}(t) = 0. \end{aligned}$$

For any $(x, y) \in \mathcal{M}^2$ and $\lambda > 0$, we also write the spectral function

$$e_{[0,\lambda]}(x, y) = \sum_{\lambda_j \in \lambda} \phi_j(x)\phi_j(y).$$

We now decompose the derivatives of $e_{[0,\lambda]}(x, y)$ via a coordinate patch $\tau : U \subset \mathbb{R}^d \rightarrow V \subset \mathcal{M}$ as follows:

$$(119) \quad \partial_x^I \partial_y^J e_{[0,\lambda]}(x, y) = \rho \star \partial_x^I \partial_y^J e_{[0,\lambda]}(x, y) + [\partial_x^I \partial_y^J e_{[0,\lambda]}(x, y) - \rho \star \partial_x^I \partial_y^J e_{[0,\lambda]}(x, y)],$$

where the convolution has to be understood with respect to λ . We now invoke the part of a work of Canzani-Hanin that holds true without any geometric property on the manifold. In local coordinates, Lemma 5 of [CH18] (with $Q = \text{Id}$, $D_0^Q = 1$ and $D_{-1}^Q = 0$) implies the following statement.

PROPOSITION 32 (Canzani-Hanin). — *Let \mathcal{M} be a boundaryless compact Riemannian manifold of dimension $d > 2$ and let us consider a coordinate patch $\tau : U \subset \mathbb{R}^d \rightarrow V \subset \mathcal{M}$ with $\text{diam}(V) \in \frac{1}{2} \text{inj}(\mathcal{M})$. For any multi-indexes $I \in \mathbb{N}^d$ and $J \in \mathbb{N}^d$, the following asymptotics holds true for any $(x, y) \in V^2$ and any $\nu > 1$ (where the spatial derivatives ∂_x and ∂_y are seen in the coordinate patch and ∂_ν is seen in \mathbb{R}):*

$$\partial_\nu(\rho \star \partial_x^I \partial_y^J e_{[0,\nu]})(x, y) = \frac{\nu^{d-1}}{(2\pi)^d} \partial_x^I \partial_y^J \left\{ \int_{|\xi|_y=1} e^{i\nu \langle \exp_y^{-1}(x), \xi \rangle} ds(\xi) \right\} + \mathcal{O}(\nu^{d-2+|I|+|J|} \delta_g(x, y) + (1 + \nu)^{d-3+|I|+|J|}),$$

where ds denotes the measure of the unit sphere of $T_y^* \mathcal{M}$ (canonically induced by the inner product on the cotangent space $T_y^* \mathcal{M}$). Finally, the remainder is uniform provided that x and y run over a compact subset of the open set V .

The presence of the distance $\delta_g(x, y)$ in the remainder is of interest in [CH18] and also in [CH15b, Prop. 10] because $\delta_g(x, y)$ may become very small in the previous papers. However, by using the fact that the Riemannian distance is bounded on the compact manifold \mathcal{M} , we may integrate on $\nu \in [0, \lambda]$ with polar coordinates to get

$$\begin{aligned} \rho \star \partial_x^I \partial_y^J e_{[0,\lambda]}(x, y) &= \frac{\partial_x^I \partial_y^J}{(2\pi)^d} \left\{ \int_0^\lambda \left(\int_{|\xi|_y=1} e^{i(\exp_y^{-1}(x), \nu\xi)} \nu^{d-1} ds(\xi) \right) d\nu \right\} + \mathcal{O}(\lambda^{d-1+|I|+|J|}) \\ &= \frac{\partial_x^I \partial_y^J}{(2\pi)^d} \left\{ \int_{|\xi|_y < \lambda} e^{i(\exp_y^{-1}(x), \xi)} \frac{d\xi}{\sqrt{|g_y|}} \right\} + \mathcal{O}(\lambda^{d-1+|I|+|J|}), \end{aligned}$$

where $d\xi/\sqrt{|g_y|}$ is the usual notation for the Lebesgue measure associated to the inner product of the cotangent space $T_y^* \mathcal{M}$. With the formulas (117), (56) and [CH18, lines (8–9)], we recover the principal term of the statement of Theorem 11:

$$\rho \star \partial_x^I \partial_y^J e_{[0,\lambda]}(x, y) = \frac{\partial_x^I \partial_y^J}{(2\pi)^{d/2}} \left\{ \int_0^\lambda \nu^{d-1} \mathfrak{d}_{d-1}(\nu \delta_g(x, y)) d\nu \right\} + \mathcal{O}(\lambda^{d-1+|I|+|J|}).$$

To complete the proof of Theorem 11, we merely need to look at the decomposition (119) and the following result (still written in local coordinates).

PROPOSITION 33. — *Let \mathcal{M} be a boundaryless compact Riemannian manifold of dimension $d > 2$ and let us consider a coordinate patch $\tau : U \subset \mathbb{R}^d \rightarrow V \subset \mathcal{M}$. For any two multi-indices $I \in \mathbb{N}^d$ and $J \in \mathbb{N}^d$, the following inequalities hold true for any x and y belonging to a compact subset of the open set V :*

$$(120) \quad \forall \lambda > 1, \quad |\rho \star \partial_x^I \partial_y^J e_{[0, \lambda]}(x, y) - \partial_x^I \partial_y^J e_{[0, \lambda]}(x, y)| \leq C \lambda^{d-1+|I|+|J|}.$$

Proof. — Let us convene that $e_{[0, \lambda]}(x, y)$ equals 0 for $\lambda < 0$. The Fourier transform of $\lambda \in \mathbb{R} \mapsto \rho \star \partial_x^I \partial_y^J e_{[0, \lambda]}(x, y) - \partial_x^I \partial_y^J e_{[0, \lambda]}(x, y)$ can be factorized by $\widehat{\rho}(t) - 1$ and hence vanishes near 0 thanks to (118). Hence, a standard Tauberian lemma (see [Sog17, Lem. 4.2.3, lines (4.2.13) and (4.2.14)]) ensures that (120) will be a consequence of the following estimates for all $\lambda > 0$ and $s \in [0, 1]$

$$|\rho \star \partial_x^I \partial_y^J e_{(\lambda, \lambda+s]}(x, y) - \partial_x^I \partial_y^J e_{(\lambda, \lambda+s]}(x, y)| \leq C(1 + \lambda)^{d-1+|I|+|J|}.$$

Clearly, it is sufficient to prove the following two inequalities

$$(121) \quad |\partial_x^I \partial_y^J e_{(\lambda, \lambda+s]}(x, y)| \leq C(1 + \lambda)^{d-1+|I|+|J|},$$

$$(122) \quad |\rho \star \partial_x^I \partial_y^J e_{(\lambda, \lambda+s]}(x, y)| \leq C(1 + \lambda)^{d-1+|I|+|J|}.$$

Let us prove (121). The Cauchy-Schwarz inequality gives

$$\begin{aligned} |\partial_x^I \partial_y^J e_{(\lambda, \lambda+s]}(x, y)| &= \left| \sum_{\lambda < \lambda_j \leq \lambda+s} \partial_x^I \phi_j(x) \partial_y^J \phi_j(y) \right| \\ &\leq \sqrt{\sum_{\lambda < \lambda_j \leq \lambda+s} |\partial_x^I \phi_j(x)|^2} \sqrt{\sum_{\lambda < \lambda_j \leq \lambda+s} |\partial_y^J \phi_j(y)|^2}. \end{aligned}$$

We then get (121) by invoking the following Bernstein-type inequality proved by Bin (see [Bin04, line (11)]):

$$(123) \quad \sqrt{\sum_{\lambda < \lambda_j \leq \lambda+1} |\partial_x^I \phi_j(x)|^2} \leq C(1 + \lambda)^{(d-1)/2+|I|}.$$

Note actually that the results of [Bin04] are stated in specific coordinate patches that are usually called normal coordinates. Here are some details explaining why (123) is still true for other coordinates. Let us consider normal coordinates

$$\tau_{\mathcal{N}} : U_{\mathcal{N}} \subset \mathbb{R}^d \rightarrow V \subset \mathcal{M}$$

(see [Bin04] for definitions). Since $\tau_{\mathcal{N}}^{-1} \circ \tau : U \rightarrow U_{\mathcal{N}}$ is a diffeomorphism, all its derivatives are bounded on any compact subset of U . Given an eigenfunction $\phi_j : \mathcal{M} \rightarrow \mathbb{R}$, it is clear that bounding a finite number of derivatives of $\phi_j \circ \tau : U \rightarrow \mathbb{R}$ or $\phi_j \circ \tau_{\mathcal{N}} : U_{\mathcal{N}} \rightarrow \mathbb{R}$ are equivalent problems since, by decomposing $\phi_j \circ \tau = (\phi_j \circ \tau_{\mathcal{N}}) \circ (\tau_{\mathcal{N}}^{-1} \circ \tau)$ and using the Faà di Bruno's formula, we have for any $k \in \mathbb{N}$ and $x \in V$ an inequality of the form

$$\sum_{|I| \leq k} (\partial^I (\phi_j \circ \tau)|_{\tau^{-1}(x)})^2 \leq C \sum_{|I| \leq k} (\partial^I (\phi_j \circ \tau_{\mathcal{N}})|_{\tau_{\mathcal{N}}^{-1}(x)})^2$$

and the last inequality is moreover uniform for x running over a compact subset of V . Hence the inequality (123) of [Bin04] still holds for non-normal coordinates.

Let us turn to the proof of (122). We begin by writing the finite sums

$$\begin{aligned} \rho \star \partial_x^I \partial_y^J e_{(\lambda, \lambda+s]}(x, y) &= \rho \star \left(\sum_{j \in \mathbb{N}} \partial_x^I \phi_j(x) \partial_y^J \phi_j(y) \mathbf{1}_{[\lambda_j-s, \lambda_j]}(\lambda) \right) \\ &= \sum_{j \in \mathbb{N}} \partial_x^I \phi_j(x) \partial_y^J \phi_j(y) \int_{\lambda-\lambda_j}^{\lambda-\lambda_j+s} \rho(\nu) d\nu. \end{aligned}$$

Using that ρ belongs to the Schwartz space and the fact that the length of the interval $[\lambda - \lambda_j, \lambda - \lambda_j + s]$ is less or equal to 1, we obtain the following upper bound

$$\begin{aligned} |\rho \star \partial_x^I \partial_y^J e_{(\lambda, \lambda+s]}(x, y)| &\leq C \sum_{j \in \mathbb{N}} \frac{|\partial_x^I \phi_j(x)| |\partial_y^J \phi_j(y)|}{(1 + |\lambda - \lambda_j|)^{d+1+|I|+|J|}} \\ &\leq C \sqrt{\sum_{j \in \mathbb{N}} \frac{|\partial_x^I \phi_j(x)|^2}{(1 + |\lambda - \lambda_j|)^{d+1+2|I|}}} \sqrt{\sum_{j \in \mathbb{N}} \frac{|\partial_y^J \phi_j(y)|^2}{(1 + |\lambda - \lambda_j|)^{d+1+2|J|}}}. \end{aligned}$$

We finish by bounding the first square root (the other is obviously similar):

$$\begin{aligned} \sum_{j \in \mathbb{N}} \frac{|\partial_x^I \phi_j(x)|^2}{(1 + |\lambda - \lambda_j|)^{d+1+2|I|}} &= \sum_{k \in \mathbb{N}} \sum_{|\lambda_j - \lambda| \in [k, k+1)} \frac{|\partial_x^I \phi_j(x)|^2}{(1 + |\lambda - \lambda_j|)^{d+1+2|I|}} \\ &\leq C \sum_{k \in \mathbb{N}} \frac{1}{(1 + k)^{d+1+2|I|}} \sum_{|\lambda_j - \lambda| \in [k, k+1)} |\partial_x^I \phi_j(x)|^2. \end{aligned}$$

In the last sum, λ_j belongs to $[\lambda - k - 1, \lambda - k] \cup [\lambda + k, \lambda + k + 1]$ and we may use (123) to get

$$\begin{aligned} \sum_{j \in \mathbb{N}} \frac{|\partial_x^I \phi_j(x)|^2}{(1 + |\lambda - \lambda_j|)^{d+1+2|I|}} &\leq C \sum_{k \in \mathbb{N}} \frac{(1 + \lambda + k)^{d-1+2|I|}}{(1 + k)^{d+1+2|I|}} \\ &\leq C(1 + \lambda)^{d-1+2|I|}. \end{aligned}$$

REFERENCES

[AT08] A. AYACHE & N. TZVETKOV – “ L^p properties for Gaussian random series”, *Trans. Amer. Math. Soc.* **360** (2008), no. 8, p. 4425–4439.
 [Bin04] X. BIN – “Derivatives of the spectral function and Sobolev norms of eigenfunctions on a closed Riemannian manifold”, *Ann. Global Anal. Geom.* **26** (2004), no. 3, p. 231–252.
 [BGT04] N. BURQ, P. GÉRARD & N. TZVETKOV – “Strichartz inequalities and the nonlinear Schrödinger equation on compact manifolds”, *Amer. J. Math.* **126** (2004), no. 3, p. 569–605.
 [BL13] N. BURQ & G. LEBEAU – “Injections de Sobolev probabilistes et applications”, *Ann. Sci. École Norm. Sup. (4)* **46** (2013), no. 6, p. 917–962.
 [BL14] ———, “Probabilistic Sobolev embeddings, applications to eigenfunctions estimates”, in *Geometric and spectral analysis*, Contemp. Math., vol. 630, American Mathematical Society, Providence, RI, 2014, p. 307–318.
 [CH15a] Y. CANZANI & B. HANIN – “Fixed frequency eigenfunction immersions and supremum norms of random waves”, *Electron. Res. Announc. Math. Sci.* **22** (2015), p. 76–86.
 [CH15b] ———, “Scaling limit for the kernel of the spectral projector and remainder estimates in the pointwise Weyl law”, *Anal. PDE* **8** (2015), no. 7, p. 1707–1731.

- [CH18] ———, “ C^∞ scaling asymptotics for the spectral projector of the Laplacian”, *J. Geom. Anal.* **28** (2018), no. 1, p. 111–122.
- [CE08] J. CHEEGER & D. G. EBIN – *Comparison theorems in Riemannian geometry*, AMS Chelsea Publishing, Providence, RI, 2008, Revised reprint of the 1975 original.
- [Die72] J. DIEUDONNÉ – *Treatise on analysis. Vol. III*, Pure and Applied Math., vol. 10-III, Academic Press, New York-London, 1972.
- [Dud67] R. M. DUDLEY – “The sizes of compact subsets of Hilbert space and continuity of Gaussian processes”, *J. Functional Analysis* **1** (1967), p. 290–330.
- [Dud14] ———, *Uniform central limit theorems*, 2nd ed., Cambridge Studies in Advanced Mathematics, vol. 142, Cambridge University Press, New York, 2014.
- [Far08] J. FARAUT – *Analysis on Lie groups. An introduction*, Cambridge Studies in Advanced Math., vol. 110, Cambridge University Press, Cambridge, 2008.
- [FTR67] A. FIGÀ-TALAMANCA & D. RIDER – “A theorem on random Fourier series on noncommutative groups”, *Pacific J. Math.* **21** (1967), no. 3, p. 487–492.
- [Gri10] S. GRIVAUX – “Almost sure convergence of some random series”, *C. R. Acad. Sci. Paris* **348** (2010), no. 3-4, p. 155–159.
- [Hör68] L. HÖRMANDER – “The spectral function of an elliptic operator”, *Acta Math.* **121** (1968), p. 193–218.
- [Ime18] R. IMEKRAZ – “Concentration et randomisation universelle de sous-espaces propres”, *Anal. PDE* **11** (2018), no. 2, p. 263–350.
- [Ime19] ———, “Multidimensional Paley-Zygmund theorems and sharp L^p estimates for some elliptic operators”, *Ann. Inst. Fourier (Grenoble)* **69** (2019), no. 6, p. 2723–2809.
- [IRT16] R. IMEKRAZ, D. ROBERT & L. THOMANN – “On random Hermite series”, *Trans. Amer. Math. Soc.* **368** (2016), no. 4, p. 2763–2792.
- [Kah85] J.-P. KAHANE – *Some random series of functions*, 2nd ed., Cambridge Studies in Advanced Math., vol. 5, Cambridge University Press, Cambridge, 1985.
- [LT91] M. LEDOUX & M. TALAGRAND – *Probability in Banach spaces. Isoperimetry and processes*, *Ergeb. Math. Grenzgeb. (3)*, vol. 23, Springer-Verlag, Berlin, 1991.
- [Ler10] N. LERNER – *Metrics on the phase space and non-selfadjoint pseudo-differential operators*, *Pseudo-Differential Operators. Theory and Appl.*, vol. 3, Birkhäuser Verlag, Basel, 2010.
- [LQ18a] D. LI & H. QUEFFÉLEC – *Introduction to Banach spaces: analysis and probability. Vol. 1*, Cambridge Studies in Advanced Math., vol. 166, Cambridge University Press, Cambridge, 2018.
- [LQ18b] ———, *Introduction to Banach spaces: analysis and probability. Vol. 2*, Cambridge Studies in Advanced Math., vol. 167, Cambridge University Press, Cambridge, 2018.
- [LT79] J. LINDENSTRAUSS & L. TZAFIRI – *Classical Banach spaces. II*, *Ergeb. Math. Grenzgeb. (3)*, vol. 97, Springer-Verlag, Berlin-New York, 1979, Function spaces.
- [Mar73] M. B. MARCUS – “Continuity of Gaussian processes and random Fourier series”, *Ann. Probab.* **1** (1973), no. 6, p. 968–981.
- [Mar75] ———, “Uniform convergence of random Fourier series”, *Ark. Mat.* **13** (1975), no. 1, p. 107–122.
- [MP81] M. B. MARCUS & G. PISIER – *Random Fourier series with applications to harmonic analysis*, *Annals of Math. Studies*, vol. 101, Princeton University Press, Princeton, NJ, 1981.
- [MS70] M. B. MARCUS & L. A. SHEPP – “Continuity of Gaussian processes”, *Trans. Amer. Math. Soc.* **151** (1970), p. 377–391.
- [Mau74] B. MAUREY – “Type et cotype dans les espaces munis de structures locales inconditionnelles”, in *Espaces L^p , applications radonifiantes et géométrie des espaces de Banach*, Séminaire Maurey-Schwartz, Ecole Polytechnique, 1974, Exp. nos. 24 & 25.
- [PZ30] R. PALEY & A. ZYGMUND – “On some series of functions. I, II”, *Proc. Camb. Phil. Soc.* **26** (1930), p. 337–357, 458–474.
- [Pis78] G. PISIER – “Sur l'espace de Banach des séries de Fourier aléatoires presque sûrement continues”, in *Séminaire sur la géométrie des espaces de Banach (1977–1978)*, École Polytechnique, Palaiseau, 1978, Exp. no. 17–18, 33p.
- [Pis89] ———, *The volume of convex bodies and Banach space geometry*, Cambridge Tracts in Math., vol. 94, Cambridge University Press, Cambridge, 1989.

- [Saf88] Y. G. SAFAROV – “Asymptotic of the spectral function of a positive elliptic operator without the nontrap condition”, *Funct. Anal. Appl.* **22** (1988), no. 3, p. 213–223.
- [SZ54] R. SALEM & A. ZYGMUND – “Some properties of trigonometric series whose terms have random signs”, *Acta Math.* **91** (1954), no. 1, p. 245–301.
- [Sch42] I. J. SCHOENBERG – “Positive definite functions on spheres”, *Duke Math. J.* **9** (1942), p. 96–108.
- [Shu01] M. A. SHUBIN – *Pseudodifferential operators and spectral theory*, 2nd ed., Springer-Verlag, Berlin, 2001.
- [Sog17] C. D. SOGGE – *Fourier integrals in classical analysis*, 2nd ed., Cambridge Tracts in Math., vol. 210, Cambridge University Press, Cambridge, 2017.
- [SW71] E. M. STEIN & G. WEISS – *Introduction to Fourier analysis on Euclidean spaces*, Princeton Math. Series, vol. 32, Princeton University Press, Princeton, NJ, 1971.
- [Sze75] G. SZEGŐ – *Orthogonal polynomials*, 4th ed., Amer. Math. Soc. Colloq. Publ., vol. XXIII, American Mathematical Society, Providence, RI, 1975.
- [Tzv10] N. TZVETROV – “Riemannian analogue of a Paley-Zygmund theorem”, in *Séminaire Équations aux Dérivées Partielles (2008–2009)*, École Polytechnique, Palaiseau, 2010, Exp. no. XV, 14p.
- [Zel09] S. ZELDITCH – “Real and complex zeros of Riemannian random waves”, in *Spectral analysis in geometry and number theory*, Contemp. Math., vol. 484, American Mathematical Society, Providence, RI, 2009, p. 321–342.

Manuscript received 12th August 2021

accepted 11th April 2022

RAFIK IMEKRAZ, La Rochelle Université, MIA, EA 3165

F-17031 La Rochelle, France

E-mail : rafik.imekraz@univ-lr.fr

Url : <https://pageperso.univ-lr.fr/rafik.imekraz/>